

OPEN GROMOV-WITTEN INVARIANTS AND SEIDEL REPRESENTATIONS FOR TORIC MANIFOLDS

KWOKWAI CHAN, SIU-CHEONG LAU, NAICHUNG CONAN LEUNG, AND HSIAN-HUA TSENG

ABSTRACT. Let X be a compact toric manifold whose anti-canonical divisor $-K_X$ is numerically effective. Let $L \subset X$ be a regular fiber of the moment map of the torus action on X . Fukaya-Oh-Ohta-Ono [14] defined open Gromov-Witten invariants of X as virtual counts of holomorphic discs with Lagrangian boundary condition L . We prove a formula for such open Gromov-Witten invariants which equates them with closed Gromov-Witten invariants of certain X -bundles over $\mathbb{P}^1$ that appear in the construction of the Seidel representations [32, 30] for X . We apply this formula and properties of Seidel representations to explicitly calculate all these open Gromov-Witten invariants. This yields a formula for the disc potential of X , and a description of the inverse of the ring isomorphism of Fukaya-Oh-Ohta-Ono [11].

CONTENTS

1. Introduction	2
1.1. Statements of results	2
1.2. Outline of methods	4
Acknowledgement	7
2. A brief review on open Gromov-Witten invariants of toric manifolds	7
3. Hori-Vafa superpotential and the toric mirror theorem	11
3.1. Mirror theorems	11
3.2. Extended moduli	14
3.3. Batyrev elements	17
4. Seidel representations for toric manifolds	20
5. Relating open and closed invariants	23
5.1. First step	24
5.2. Second step	31
6. Computing closed invariants by Seidel representations	33
6.1. Calculations	33
6.2. Corollaries	36
References	38

1. INTRODUCTION

1.1. Statements of results. Let X be a complex n -dimensional compact toric manifold equipped with a toric Kähler form ω . X admits a Hamiltonian action by a complex torus $\mathbf{T}_{\mathbb{C}} \simeq (\mathbb{C}^*)^n$. Let $L \subset X$ be a regular fiber of the associated moment map. Then L is diffeomorphic to $(S^1)^n$ and is a Lagrangian submanifold of X . We will refer to it as a Lagrangian torus fiber. Let $\beta \in \pi_2(X, L)$ be a relative homotopy class represented by a disc bounded by L . In [14], Fukaya-Oh-Ohta-Ono defined the genus-zero *open Gromov-Witten invariant* $n_1(\beta) \in \mathbb{Q}$ as a virtual count of holomorphic discs in X bounded by L representing the class β . The precise definition of $n_1(\beta)$ is reviewed in Definition 2.1. Genus-zero open Gromov-Witten invariants assemble to a generating function W^{LF} called the *disc potential*, see Definition 2.4.

The disc potential W^{LF} plays a fundamental role in the symplectic geometry of the toric manifold X . It was used by Fukaya-Oh-Ono-Ohta [14, 15, 11] to detect non-displaceable Lagrangian torus fibers in X . Moreover, the A_{∞} algebra, which encodes all symplectic information of a Lagrangian torus fiber, is determined by W^{LF} by taking its derivatives. In an upcoming work of Abouzaid-Fukaya-Oh-Ohta-Ono, it is shown that the Fukaya category of X is indeed generated by Lagrangian torus fibers. As a consequence, W^{LF} completely determines the Fukaya category of X . W^{LF} is also important in the study of mirror symmetry for toric manifolds because it serves as the Landau-Ginzburg mirror of X and its Jacobian ring determines the quantum cohomology of X [11].

Open Gromov-Witten invariants are in general very difficult to compute, especially when the obstruction of the corresponding moduli space is non-trivial. For Fano toric manifolds where the obstruction bundle is trivial, open Gromov-Witten invariants were computed by Cho-Oh [10]. The next simplest non-trivial example, which is the Hirzebruch surface $\mathbb{F}_2$, was computed by Auroux [2] using wall-crossing technique¹ and by Fukaya-Oh-Ohta-Ono [16] using degenerations. Later, under certain restrictions on the geometry of the toric manifolds, open Gromov-Witten invariants were computed in [4, 6, 24, 5, 8]. On the other hand, assuming convergence of W^{LF} , it was proved in [7] that open Gromov-Witten invariants of semi-Fano toric manifolds can be extracted from the inverse mirror maps.

One purpose of this paper is to compute open Gromov-Witten invariants $n_1(\beta)$ for *all compact semi-Fano toric manifolds*. By definition, a toric manifold X is semi-Fano if we have $-K_X \cdot C \geq 0$ for every holomorphic curve $C \subset X$. Let $\beta \in \pi_2(X, L)$ be a disc class² of Maslov index two such that $n_1(\beta) \neq 0$. By the results of Cho-Oh [10] and Fukaya-Oh-Ohta-Ono [14] (see also Lemma 2.3), β must be of the form $\beta = \beta_l + \alpha$ where $\beta_l \in \pi_2(X, L)$ is the *basic disc class* associated to a toric prime divisor D_l and $\alpha \in H_2(X, \mathbb{Z})$ is an effective curve class with $-K_X \cdot \alpha = 0$. Define the following generating function

$$\delta_l(q) := \sum_{\substack{\alpha \in H_2(X, \mathbb{Z}) \setminus \{0\} \text{ effective} \\ -K_X \cdot \alpha = 0}} n_1(\beta_l + \alpha) q^{\alpha},$$

¹Using the same techniques, Auroux [2] also computed the open Gromov-Witten invariants for the Hirzebruch surface $\mathbb{F}_3$.

²By dimension reasons only classes β of Maslov index two can have non-zero $n_1(\beta)$. See Section 2 for a discussion on Maslov index.

see Definition 2.4 for more details.

One of our main results is an explicit formula for the generating function $\delta_l(q)$, which may be stated as follows. The so-called *toric mirror theorem* of Givental [18] and Lian-Liu-Yau [27], as recalled in Theorem 3.4, states that there is an equality

$$I(\check{q}, z) = J(q(\check{q}), z),$$

where $I(\check{q}, z)$ is the combinatorially defined *I-function* of X (see Definition 3.1), $J(q, z)$ is a certain generating function of closed Gromov-Witten invariants of X called the *J-function* (see equation (3.3)), and $q(\check{q})$ is the *mirror map* in Definition 3.2. Our formula for $\delta_l(q)$ reads as follows:

Theorem 1.1. *Let X be a compact semi-Fano toric manifold. Then*

$$1 + \delta_l(q) = \exp(g_l(\check{q}(q))),$$

where

$$g_l(\check{q}) := \sum_d \frac{(-1)^{(D_l \cdot d)} (-(D_l \cdot d) - 1)!}{\prod_{p \neq l} (D_p \cdot d)!} \check{q}^d$$

where the summation is over all curve classes d satisfying

$$-K_X \cdot d = 0, D_l \cdot d < 0 \text{ and } D_p \cdot d \geq 0 \text{ for all } p \neq l$$

and $\check{q} = \check{q}(q)$ is the inverse of the mirror map $q = q(\check{q})$.

The toric mirror map $q = q(\check{q})$ is combinatorially explicitly defined, and its inverse $\check{q} = \check{q}(q)$ can be explicitly computed, at least recursively. Therefore our formula provides an effective calculation for all genus-zero open Gromov-Witten invariants.

Our formula may be inverted to give a formula which expresses the inverse mirror map $\check{q}(q)$ in terms of open Gromov-Witten invariants, see Corollary 6.7. This gives the inverse mirror map an enumerative meaning in terms of disc counting.

Our calculation of open Gromov-Witten invariants can also be neatly stated in terms of the disc potential:

Theorem 1.2. *Let X be a compact semi-Fano toric manifold. Then*

$$(1.1) \quad W_q^{\text{LF}} = \tilde{W}_{\check{q}(q)}^{\text{HV}},$$

where $\tilde{W}^{\text{HV}}$ is the Hori-Vafa superpotential for X in a certain explicit choice of coordinates of $(\mathbb{C}^*)^n$, see Definition 3.5 and Equation (3.4).

Theorem 1.2 may be considered as an *open mirror theorem* for toric manifolds. From this open mirror theorem, the disc potential W^{LF} is exactly the same as the Hori-Vafa superpotential $\tilde{W}^{\text{HV}}$. While the disc potential W^{LF} is a relatively new object invented to describe the symplectic geometry of X , the Hori-Vafa superpotential $\tilde{W}^{\text{HV}}$ has been studied extensively in the literature. Thus existing knowledge on $\tilde{W}^{\text{HV}}$ can be used to understand the disc potential W^{LF} better. In particular, since $\tilde{W}^{\text{HV}}$ is written in terms of (inverse) mirror maps which are known to be convergent, it follows that

Theorem 1.3. *The coefficients of the disc potential W^{LF} of a compact semi-Fano toric manifold X are convergent power series in the Kähler parameters q .*

Moreover we have the mirror theorem [18, 27] which gives an isomorphism

$$(1.2) \quad QH^*(X, \omega_q) \xrightarrow{\simeq} \text{Jac}(\tilde{W}^{HV})$$

between the quantum cohomology of X and the Jacobian ring of the Hori-Vafa superpotential when X is semi-Fano. Combining with Theorem 1.2, this gives another proof that

Corollary 1.4 (FOOO's isomorphism [11] for small quantum cohomology in semi-Fano case³). *Let X be a compact semi-Fano toric manifold. Then there exists an isomorphism*

$$(1.3) \quad QH^*(X, \omega_q) \xrightarrow{\simeq} \text{Jac}(W_q^{LF})$$

between the small quantum cohomology ring of X and the Jacobian ring of the disc potential W_q^{LF} .

On the other hand, Seidel [32] (for monotone cases) and McDuff-Tolman [31] constructed an isomorphism in the reverse direction by using Seidel representations, which were in fact also used by Fukaya-Oh-Ohta-Ono [11] in their proof of the injectivity of the homomorphism (1.3) (without knowing their explicit relations). Using our results on open Gromov-Witten invariants, we deduce that they are indeed inverse to each other:

Theorem 1.5. *Suppose X is semi-Fano. Then the isomorphism (1.3) maps the (normalized) Seidel elements $S_l^\circ \in QH^*(X, \omega_q)$ (see Section 4) to the generators $Z_1, \dots, Z_m$ of the Jacobian ring $\text{Jac}(W_q^{LF})$, where Z_l are monomials defined by Equation (2.2).*

We conjecture that Theorem 1.5, which provides a highly non-trivial relation between open Gromov-Witten invariants and Seidel representations, holds true for *all* toric manifolds. See Conjecture 6.12 for the precise statement.

1.2. Outline of methods. The *closed* Gromov-Witten theory for toric manifolds has been studied extensively, and various computational tools such as virtual localization are available and very powerful. The situation is drastically different for *open* Gromov-Witten theory with respect to Lagrangian torus fibers: the open Gromov-Witten invariants, which are defined using the geometry and analysis of moduli spaces of pseudo-holomorphic discs whose structures are very sophisticated, are very hard to compute, especially because of the lack of localization techniques⁴.

In our previous paper [7] we took an analytic approach to study open Gromov-Witten invariants of toric manifolds. In [7, Theorem 1.2] we proved Theorem 1.2 under the technical assumption that the generating functions $\delta_l(q)$ are convergent power series in q . Our proof employs the theory of universal unfolding of analytic singularities and uses the isomorphism $\text{Jac}(W_q^{LF}) \simeq \text{Jac}(W_{\tilde{q}(q)}^{HV})$ deduced from the toric mirror theorem of Givental [18] and Lian-Liu-Yau [27] and the isomorphism (1.3).

³Fukaya-Oh-Ohta-Ono proved a ring isomorphism between the *big* quantum cohomology ring of a compact toric manifold X and the Jacobian ring of the so-called *bulk-deformed* potential function. Our results deduces their isomorphism for the small quantum cohomology of a semi-Fano toric manifold X .

⁴The situation is rather different for open Gromov-Witten invariants of toric Calabi-Yau threefolds with respect to Aganagic-Vafa type Lagrangian submanifolds—in this case the invariants are practically defined by localization formula and can certainly be evaluated using it.

In this paper we study the problem of computing open Gromov-Witten invariants via a geometric approach entirely different from the one in [7]. Let us briefly explain the ideas here. By the classification result of Cho-Oh [10], a stable disc of Maslov index two in a toric manifold X must have its domain being a union of a disc D and a collection of rational curves. Naïvely one may hope to “cap off” the disc D by finding another disc D' and gluing D and D' together along their boundaries to form a sphere. If this can be done, it is then natural to speculate that the open Gromov-Witten invariants we want to compute are equal to certain closed Gromov-Witten invariants. This idea was first successfully worked out by Chan in [4] for toric manifolds of the form $X = \mathbb{P}(K_Y \oplus \mathcal{O}_Y)$ where Y is a compact Fano toric manifold. In this case the $\mathbb{P}^1$ -bundle structure on X provides a way to find the needed disc D' . This idea was also applied in subsequent works by Lau-Leung-Wu [23, 24], Chan-Lau-Leung [6], Chan-Lau [5] and Chan-Lau-Tseng [8]. Through these works, it became clear that in more general situations, in order to find the “capping-off” disc D' , one may need to work with a toric target space different from X .

Here we further advance this geometric approach to determine open Gromov-Witten invariants for *all* compact semi-Fano toric manifolds. One novelty of this paper is the use of *Seidel spaces*. Given a toric manifold X , let D_l be a toric prime divisor corresponding to a ray in the fan and let v_l be the generator of this ray. Then $-v_l$ defines a $\mathbb{C}^*$ -action on X . Let $\mathbb{C}^*$ act on $\mathbb{C}^2 \setminus \{0\}$ by $z \cdot (u, v) := (zu, zv)$, $z \in \mathbb{C}^*$, $(u, v) \in \mathbb{C}^2 \setminus \{0\}$. The *Seidel space* associated to the $\mathbb{C}^*$ -action defined by $-v_l$ is the quotient

$$E_l^- := (X \times (\mathbb{C}^2 \setminus \{0\})) / \mathbb{C}^*.$$

By construction, E_l^- is also a toric manifold. Moreover there is a natural map $E_l^- \rightarrow \mathbb{P}^1$ which gives E_l^- the structure of a fiber bundle whose fibers are diffeomorphic to X . The toric data of E_l^- , as well as geometric information such as the Mori cone, can be explicitly described. See Section 4 for more details.

Recall that the disc classes which give non-zero open Gromov-Witten invariants are of the form $\beta_l + \alpha$, where β_l is the basic disc class associated to the toric prime divisor D_l for some l , and $\alpha \in H_2(X, \mathbb{Z})$ is an effective curve class with $-K_X \cdot \alpha = 0$. We prove the following

Theorem 1.6 (See Theorem 5.1). *Let X be a compact semi-Fano toric manifold and $L \subset X$ a Lagrangian torus fiber. Then for any toric prime divisor D_k , we have*

$$(1.4) \quad n_{1,1}^X(\beta_l + \alpha; D_k; [\text{pt}]_L) = \langle D_k^{E_l^-}, [\text{pt}]_{E_l^-} \rangle_{0,2,\sigma_l^- + \alpha}^{E_l^-}.$$

On the left-hand side of (1.4), $n_{1,1}^X(\beta_l + \alpha; D_k; [\text{pt}]_L)$ is the open Gromov-Witten invariant defined in Definition 2.1, which roughly speaking counts discs of classes $\beta_l + \alpha$ meeting a fixed point in L at the boundary marked point and meeting the divisor D_k at the interior marked point. The invariant $n_{1,1}^X(\beta_l + \alpha; D_k; [\text{pt}]_L)$ is related to the previous invariant $n_1(\beta_l + \alpha)$ via the *divisor equation* proved by Cho [9]:

$$n_{1,1}^X(\beta_l + \alpha; D_k; [\text{pt}]_L) = (D_k \cdot (\beta_l + \alpha)) n_1(\beta_l + \alpha),$$

see Theorem 2.2.

On the right-hand side of (1.4) we have the two-point *closed* Gromov-Witten invariant

$$(1.5) \quad \langle D_k^{E_l^-}, [\text{pt}]_{E_l^-} \rangle_{0,2,\sigma_l^- + \alpha}^{E_l^-}$$

of the Seidel space E_l^- , see Section 4 for the notations. The geometric idea behind the proof of (1.4) is the following. A stable disc representing the class $\beta_l + \alpha$ is a union of a disc Δ in X representing β_l and a rational curve C in X representing α . We identify X with the fiber of $E_l^- \rightarrow \mathbb{P}^1$ over $0 \in \mathbb{P}^1$ and consider C as a rational curve in E_l^- . The key point is that the disc Δ in X bounded by a Lagrangian torus fiber L of X can be identified with a disc $\tilde{\Delta}$ in E_l^- bounded by a Lagrangian torus fiber $\tilde{L}$ of E_l^- , and there exists a “capping-off” disc $\tilde{\Delta}'$ in E_l^- which can be glued together with $\tilde{\Delta}$ to form a rational curve representing a section σ_l^- of $E_l^- \rightarrow \mathbb{P}^1$. This idea, which is illustrated in Figure 1, allows us to identify the relevant moduli spaces. A further analysis on the Kuranishi structures yields the formula (1.4).




FIGURE 1. Relating disc invariants to Gromov-Witten invariants of the Seidel space. A disc Δ in $\mathbb{P}^1$ bounded by a torus fiber is ‘pushed’ into the associated Seidel space and compactified to a sphere in the section class σ_l^- .

Our formula (1.4) reduces the computation of open Gromov-Witten invariants to the computation of closed Gromov-Witten invariants (1.5). Such closed Gromov-Witten invariants are encoded in derivatives of the J -function of E_l^- . One may thus hope to calculate the invariants (1.5) via the J -function. Unfortunately an explicit formula for the J -function of E_l^- is not available from the toric mirror theorem in general, since E_l^- may *not* be semi-Fano.

We calculate (1.5) by a different method, which we now briefly explain. The starting point is the observation that (1.5) appears in the quantum product $S_l^- * D_k$ of the divisor D_k and the *Seidel element* S_l^- defined using the Seidel space E_l^- , see Definition 4.4. Recall that E_l^- is defined using the $\mathbb{C}^*$ -action associated to the element $-v_l$. The element v_l also defines a $\mathbb{C}^*$ -action and consequently a Seidel space E_l . Let S_l be the Seidel element defined using E_l . Since $v_l + (-v_l) = 0$, the two $\mathbb{C}^*$ -actions are inverse to each other. A basic property of Seidel elements implies that

$$S_l * S_l^- = q^{\sigma_l + \sigma_l^-},$$

see Proposition 4.5. It follows that

$$(1.6) \quad S_l * S_l^- * D_k = q^{\sigma_l + \sigma_l^-} D_k.$$

The Seidel space E_l is semi-Fano and the corresponding Seidel element S_l can be calculated using the J -function. Indeed in [20] the Seidel element S_l is explicitly calculated and expressed in terms of the so-called Batyrev elements. Their result is recalled in Proposition 6.1. We then use this calculation of S_l to extract the invariants (1.5) from (1.6). This yields a proof of Theorem 1.1. The details are given in Section 6.

Remark 1.7.

- (1) *Seidel representations were used by McDuff-Tolman [31] to derive a presentation of the quantum cohomology ring of a compact toric manifold. This presentation of the quantum cohomology ring was used in [11] to prove the isomorphism (1.3). Our approach uses the geometry of Seidel spaces directly, as well as properties of Seidel elements.*
- (2) *Theorem 1.3 gives a convergence result on the generating functions $\delta_l(q)$ of open Gromov-Witten invariants. This verifies the assumption needed in the analytic approach in our previous work [7], and thus provides an alternative way to prove Theorem 1.2 and its corollaries.*
- (3) *Equation (1.4) is a relation between open and closed Gromov-Witten invariants. An “open/closed relation” of somewhat different flavor is present in open Gromov-Witten theory of toric Calabi-Yau threefolds with respect to Aganagic-Vafa type Lagrangian branes. See [29, 25, 28, 26] for more details.*
- (4) *During the preparation of this paper, we learnt of an independent upcoming work of González-Iritani [19] in which a different approach to Theorem 1.2 is developed.*

The rest of this paper is organized as follows. Section 2 contains a brief review of open Gromov-Witten invariants of toric manifolds. In Section 3 we describe the toric mirror theorem of [18], [27], the Hori-Vafa superpotentials, the mirror maps, and related materials. In Section 4 we recall some basic materials on Seidel representations of toric manifolds. In Section 5 we prove the relation (1.4) between open and closed Gromov-Witten invariants. In Section 6 we calculate the closed Gromov-Witten invariants which appear in (1.4) and prove our main Theorems 1.1, 1.2, 1.5.

ACKNOWLEDGEMENT

We express our deep gratitude to Eduardo González and Hiroshi Iritani for various illuminating discussions on this subject and informing us another insightful approach to attack this problem. Parts of this work were carried out when the authors met at The Chinese University of Hong Kong, University of Wisconsin-Madison, Ohio State University, and Kavli Institute for the Physics and Mathematics of the Universe. We thank these institutions for their hospitality and support. The work of K. Chan described in this paper was substantially supported by a grant from the Research Grants Council of the Hong Kong Special Administrative Region, China (Project No. CUHK404412). The work of S.-C. Lau described in this paper was supported by Harvard University; he also thanks Kaoru Ono and Cheol-Hyun Cho for useful discussions on this subject. The work of N.C. Leung described in this paper was substantially supported by a grant from the Research Grants Council of the Hong Kong Special Administrative Region, China (Project No. CUHK401809).

2. A BRIEF REVIEW ON OPEN GROMOV-WITTEN INVARIANTS OF TORIC MANIFOLDS

This section aims to give a quick review on toric manifolds and their open Gromov-Witten invariants which are the central objects studied by this paper. For a detailed study of toric varieties, the readers are referred to Fulton’s excellent book [17]. The Lagrangian-Floer theory that we used in this paper is developed by Fukaya-Oh-Ohta-Ono [12, 13, 14, 15, 11].

We work with a projective toric manifold X of complex dimension n equipped with a toric Kähler form, which we assume the readers are familiar with. Let $N \cong \mathbb{Z}^n$ be a lattice. The projective toric manifold X is defined by a complete simplicial fan Σ , which is supported in the real vector space $N \otimes_{\mathbb{Z}} \mathbb{R}$. The minimal generators of rays of Σ are denoted by $v_j \in N$ for $j = 1, \dots, m$. Each v_j corresponds to a toric prime divisor denoted by $D_j \subset X$. There is an action on X by the torus $\mathbf{T}_{\mathbb{C}} = (N \otimes_{\mathbb{Z}} \mathbb{C})/N$ which preserves the Kähler structure. The moment map of this action maps X to a polytope denoted by P . Each regular fiber of the moment map is a Lagrangian submanifold, and it is a free orbit under the real torus $\mathbf{T} = (N \otimes_{\mathbb{Z}} \mathbb{R})/N$. By abuse of notation we also denote such a fiber by $\mathbf{T}$, and call it a Lagrangian torus fiber. X is said to be *semi-Fano* if $-K_X$ is numerically effective, that is, $-K_X \cdot C \geq 0$ for any holomorphic curve C . We call $-K_X \cdot C$ the Chern number of C , and also denote it by $c_1(C)$.

Let X be a semi-Fano toric manifold equipped with a toric Kähler form. Our goal is to compute the open Gromov-Witten invariants of X , which are rational numbers associated to disc classes $\beta \in \pi_2(X, \mathbf{T})$ bounded by a Lagrangian torus fiber.

We first recall the definition of open Gromov-Witten invariants. The Maslov index of a disc class β is denoted as $\mu(\beta)$. In the toric case

$$\mu(\beta) = 2 \sum_{j=1}^m (\beta \cdot D_j)$$

where $\beta \cdot D_j$ denotes the intersection number of β with the toric divisor D_j . By Cho-Oh [10], holomorphic disc classes in $\pi_2(X, \mathbf{T})$ are generated by basic disc classes β_j , $j = 1, \dots, m$, which are in one-one correspondence with the rays v_j of the fan. Since $\mu(\beta_j) = 2$ for all $j = 1, \dots, m$, every non-constant holomorphic disc has Maslov index at least two.

For each disc class $\beta \in \pi_2(X, \mathbf{T})$, Fukaya-Oh-Ohta-Ono [14, 15] defined the moduli space

$$\mathcal{M}_{l,k}^{\text{op}}(\beta)$$

of stable discs with l interior marked points and k boundary marked points representing β , which is oriented and compact. When $l = 0$, we simply denote $\mathcal{M}_{0,k}^{\text{op}}(\beta)$ by $\mathcal{M}_k^{\text{op}}(\beta)$. Here we use the superscript “op” to remind ourselves that it is the moduli space for defining open Gromov-Witten invariants. Later we will use the superscript “cl” (which stands for “closed”) for the moduli space of rational curves.

The main difficulty in defining the invariants is the lack of transversality: the actual dimension of $\mathcal{M}_{l,k}(\beta)$ in general is higher than its expected (real) dimension $n + \mu(\beta) + k + 2l - 3$. To tackle this, Fukaya-Oh-Ohta-Ono analyzed the obstruction theory and used the torus action on $\mathcal{M}_{l,k}(\beta)$ to construct a virtual fundamental chain

$$[\mathcal{M}_{l,k}(\beta)]_{\text{virt}},$$

which is a $\mathbb{Q}$ -chain of dimension $n + \mu(\beta) + k + 2l - 3$ in $X^l \times \mathbf{T}^k$. In this paper we shall only need the cases $k = 1$ and l is either 0 or 1. In these cases, when $\mu(\beta) = 2$, since non-constant stable discs bounded by $\mathbf{T}$ have Maslov indices at least two, $\mathcal{M}_{l,k}(\beta)$ has no codimension-one boundary and so $[\mathcal{M}_{l,k}(\beta)]_{\text{virt}}$ is actually a cycle. (For a nice discussion of this issue the reader is referred to [1, Section 3]). Also notice that for $k = 1$, $l = 0$ and $\mu(\beta) > 2$, $[\mathcal{M}_1(\beta)]_{\text{virt}} = 0$ due to dimension reason.

Definition 2.1 (Open Gromov-Witten invariant [14]). *Let X be a compact semi-Fano toric manifold and $\mathbf{T}$ a Lagrangian torus fiber of X . We denote by $(\cdot, \cdot)$ the Poincaré pairing on $H^*(\mathbf{T})$.*

The one-point open Gromov-Witten invariant associated to a disc class $\beta \in \pi_2(X, \mathbf{T})$ with $\mu(\beta) = 2$ is defined to be

$$n_1(\beta; [\text{pt}]_{\mathbf{T}}) = n_\beta := ([\mathcal{M}_1(\beta)]_{\text{virt}}, [\text{pt}]_{\mathbf{T}}) \in \mathbb{Q}.$$

$n_\beta := 0$ if $\mu(\beta) \neq 2$.

For a toric cycle $C \subset X$ of real codimension $\text{codim}(C)$ and a disc class $\beta \in \pi_2(X, \mathbf{T})$, let

$$\mathcal{M}_{1;1}(\beta; C) := \mathcal{M}_{1;1}(\beta) \times_X C$$

where the fiber product over X is defined by the evaluation map $\text{ev}_+ : \mathcal{M}_{1;1}(\beta) \rightarrow X$ at the interior marked point and the inclusion map $C \hookrightarrow X$. The expected dimension of $\mathcal{M}_{1;1}(\beta; C)$ is $\dim C + \mu(\beta)$. The one-point open Gromov-Witten invariant of class β relative to C is defined to be

$$n_{1,1}(\beta; C; [\text{pt}]_{\mathbf{T}}) := ([\mathcal{M}_{1;1}(\beta; C)]_{\text{virt}}, [\text{pt}]_{\mathbf{T}}) \in \mathbb{Q}$$

where the torus action on $\mathcal{M}_{1;1}(\beta; C)$ is used to construct the virtual fundamental chain $[\mathcal{M}_{1;1}(\beta; C)]_{\text{virt}}$.

Notice that by dimension counting, $n_{1,1}(\beta; C; [\text{pt}]_{\mathbf{T}}) = 0$ if $\mu(\beta) \neq \text{codim}(C)$. Thus when C is a toric divisor, we only need to consider those β with $\mu(\beta) = 2$, and the corresponding moduli $\mathcal{M}_{1;1}(\beta; C)$ has no codimension one boundary. Cho [9] has proved the following analog of divisor equation in the open case:

Theorem 2.2 (Divisor equation in the open case [9]). *For a toric divisor $D \subset X$ and a disc class $\beta \in \pi_2(X, \mathbf{T})$ with $\mu(\beta) = 2$, we have*

$$n_{1,1}(\beta; D; [\text{pt}]_{\mathbf{T}}) = (D \cdot \beta) n_1(\beta)$$

where $D \cdot \beta$ denotes the intersection number between D and β .

By using the above definition of open Gromov-Witten invariants, one can define the disc potential as follows.

First we make the following choices. By relabeling the generators $\{v_j\}_{j=1}^m$ of rays if necessary, we may assume that $v_1, \dots, v_n$ spans a cone in the fan Σ so that $\{v_1, \dots, v_n\}$ gives an integral basis of N . Denote the dual basis by $\{\nu_k\}_{k=1}^n \subset M := N^*$. Moreover, fix the basis $\{\Psi_k\}_{k=1}^{m-n}$ of $H_2(X)$ where

$$(2.1) \quad \Psi_k := - \sum_{p=1}^n (\nu_p, v_{n+k}) \beta_p + \beta_{n+k} \in H_2(X), \quad \text{for } k = 1, \dots, m-n.$$

(Recall that the basic disc classes $\{\beta_j\}_{j=1}^m$ form a basis of $H_2(X, \mathbf{T})$.) Note that $\Psi_k \in H_2(X)$ because

$$\partial \Psi_k = - \sum_{p=1}^n (\nu_p, v_{n+k}) v_p + v_{n+k} = -v_{n+k} + v_{n+k} = 0.$$

Since $(\Psi_k, D_{n+r}) = \delta_{kr}$ for all $k, r = 1, \dots, k$, the dual basis of $\{\Psi_k\}_{k=1}^{m-n}$ is $[D_{n+1}], \dots, [D_m] \in H^2(X)$.

The basis $\{\Psi_k\}_{k=1}^{m-n}$ defines flat coordinates on $H^2(X, \mathbb{C})/2\pi\mathbf{i}H^2(X, \mathbb{Z})$ by sending

$$[\eta] \in H^2(X, \mathbb{C})/2\pi\mathbf{i}H^2(X, \mathbb{Z})$$

to $q_k([\eta]) = q^{\Psi_k}(\eta) := \exp\left(\int_{\Psi_k} \eta\right)$ for $k = 1, \dots, m-n$. The complexified Kähler cone

$$\mathcal{K}_X^{\mathbb{C}} := \mathcal{K}_X \oplus (\mathbf{i}H^2(X, \mathbb{R})/2\pi\mathbf{i}H^2(X, \mathbb{Z}))$$

is embedded as an open subset of $H^2(X, \mathbb{C})/2\pi\mathbf{i}H^2(X, \mathbb{Z})$ by taking $\omega \in \mathcal{K}_X^{\mathbb{C}}$ to $-\omega \in H^2(X, \mathbb{C})/2\pi\mathbf{i}H^2(X, \mathbb{Z})$. Then $(q_k)_{k=1}^{m-n}$ is pulled back to give a flat coordinate system on $\mathcal{K}_X^{\mathbb{C}}$, so that the large radius limit has coordinate $q = 0$.

The disc potential is defined by summing the one-point open Gromov-Witten invariants for all $\beta \in \pi_2(X, \mathbf{T})$ weighted by q^β . By dimension reasons, only those β with Maslov index two contribute. Since X is a semi-Fano toric manifold, stable discs with Maslov index two must be of the form $\beta_j + \alpha$ for some basic disc class β_j and $\alpha \in H_2^{\text{eff}}(X)$ with $c_1(\alpha) = 0$. Here $H_2^{\text{eff}}(X) \subset H_2(X, \mathbb{Z})$ is the semi-group of effective curve classes of X . More precisely,

Lemma 2.3 ([10, 14]). *Let X be a semi-Fano toric manifold and $\mathbf{T}$ a Lagrangian torus fiber of X . A stable disc in $\mathcal{M}_{l,k}^{\text{op}}(\beta)$ for $k = 1$ and $l = 0, 1$ where $\beta \in \pi_2(X, \mathbf{T})$ with $\mu(\beta) = 2$ is a union of a holomorphic disc component and a rational curve, which are attached to each other at only one nodal interior point. The disc component represents the class β_j for some $j = 1, \dots, m$, and the rational curve has $c_1 = 0$. Thus the class of every stable disc is of the form $\beta_j + \alpha$ for some $j = 1, \dots, m$ and $\alpha \in H_2^{\text{eff}}(X)$ with $c_1(\alpha) = 0$.*

Proof. For a toric manifold X , by the classification theorem of Cho-Oh [10], a smooth non-constant holomorphic disc bounded by a Lagrangian torus fiber has Maslov index at least two. Moreover, a smooth holomorphic disc which has Maslov index two must represent a basic disc class β_j for some $j = 1, \dots, m$. If X is semi-Fano, every holomorphic curve has non-negative Chern number. Now a non-constant stable disc bounded by a Lagrangian torus fiber consists of at least one holomorphic disc component and possibly several sphere components. Thus it has Maslov index at least two, and if it is of Maslov index two, it must consist of only one disc component which represents a basic class β_j for some $j = 1, \dots, m$. Moreover, the sphere components have Chern number zero, and thus they are contained in the toric divisors (otherwise they are constant and cannot be stable since there is only one interior marked point). But a holomorphic disc in class β_j intersect with $\cup_k D_k$ at only one point. Thus it is only attached with one of the sphere components, and by connectedness the sphere components form a rational curve whose class is denoted as α which has $c_1 = 0$. $\square$

Moreover by the result of Cho-Oh [10], $n_1(\beta_j) = 1$ for all $j = 1, \dots, m$. Hence the disc potential is of the following form:

Definition 2.4 (Disc potential [14]). *For a semi-Fano toric manifold X , the disc potential of X is defined by*

$$W^{\text{LF}} := \sum_{l=1}^m (1 + \delta_l) Z_l$$

where

$$(2.2) \quad Z_l = \begin{cases} z_l & \text{when } l = 1, \dots, n; \\ q_{l-n} z_l & \text{when } l = n+1, \dots, m, \end{cases}$$

$$\delta_l := \sum_{\alpha \in H_2^{c_1=0}(X) - \{0\}} n_1(\beta_l + \alpha) q^\alpha$$

for all $l = 1, \dots, m$, $q^\alpha = \prod_{k=1}^{m-n} q_k^{(\alpha, D_{n+k})}$ and $H_2^{c_1=0}(X) \subset H_2^{\text{eff}}(X)$ denotes the semi-group of all effective curve classes α of X with $c_1(\alpha) = 0$. We also call W^{LF} the Lagrangian Floer superpotential of X and the superscript “LF” here stands for “Lagrangian Floer”.

Notice that each δ_l is a priori a formal power series contained in the Novikov ring in the formal variables $q_1, \dots, q_k$. One key theorem in this paper shows that W^{LF} is equal to the Hori-Vafa superpotential (up to the toric mirror map). Since by definition each coefficient of the Hori-Vafa superpotential converges, it follows that each δ_l is indeed a convergent power series.

As emphasized by Fukaya-Oh-Ohta-Ono [11], in order to extract symplectic-geometric information of the Lagrangian torus fiber from the disc potential W^{LF} , one has to restrict W^{LF} to a formal neighborhood of the large volume limit in one direction. Explicitly, we fix a Kähler class ω and substitute $q_k = T^{\int_{\Psi_k} \omega}$ for $k = 1, \dots, m-n$, where T is the formal Novikov parameter. Then δ_l becomes an element in the Novikov ring of one variable T , and W^{LF} is exactly the m_0 -term of the Lagrangian torus fiber. For toric manifolds, m_0 governs the whole Lagrangian Floer theory. All the higher A_∞ products $m_k, k \in \mathbb{N}$ can be recovered by taking the derivatives of m_0 . It can be used to detect the non-displaceable Lagrangian torus fibers. For a detailed treatment in this direction, the readers are referred to [14, 15, 11].

3. HORI-VAFA SUPERPOTENTIAL AND THE TORIC MIRROR THEOREM

After a brief review on the Lagrangian Floer theory of toric manifolds, which is the so-called A -side of mirror symmetry, we now come to the complex geometry, the so-called B -side, of the mirror. The mirror of a toric variety X is given by a Laurent polynomial W^{HV} explicitly determined by the fan of X [18, 21]. It defines a singularity theory whose moduli has flat coordinates given by the oscillatory integrals. These have explicit formulas and they will be reviewed in this section. Then we will recall the celebrated mirror theorem for toric varieties [18, 27].

3.1. Mirror theorems. The mirror complex moduli is defined as a certain neighborhood of 0 of $(\mathbb{C}^*)^{m-n}$ (see Definition 3.3), whose coordinates are denoted as $\check{q} = (\check{q}_1, \dots, \check{q}_{m-n}) \in (\mathbb{C}^*)^n$. $\check{q}_k$ is also denoted as $\check{q}^{\Psi_k}$ for $k = 1, \dots, m-n$.

The I -function is an $H^2(X, \mathbb{C})$ -valued function on the mirror complex moduli defined as follows:

Definition 3.1 (I -function). *The I -function of a toric manifold X is defined as*

$$I^X(\check{q}, z) := \exp \left(\frac{1}{z} \sum_{k=1}^{m-n} (\log \check{q}_k) [D_{n+k}] \right) \sum_{d \in H_2^{\text{eff}}(X)} \check{q}^d I_d$$

where

$$I_d := \prod_{l=1}^m \frac{\prod_{s=-\infty}^0 (D_l + sz)}{\prod_{s=-\infty}^{D_l \cdot d} (D_l + sz)}$$

and $\check{q}^d := \prod_{k=1}^{m-n} \check{q}_k^{(d, D_{n+k})}$.

From the above definition, we see that I_d can be regarded as a $\bigwedge^* H_{\mathbb{C}}^2(X, \mathbf{T})$ -valued function instead of $H^*(X, \mathbb{C})$ -valued, where $H_{\mathbb{C}}^2(X, \mathbf{T}) := H^2(X, \mathbf{T}) \otimes \mathbb{C}$. A basis of $H^2(X, \mathbf{T})$ is given by $\{D_l\}_{l=1}^m$, which is dual to the basis $\{\beta_l\}_{l=1}^m \subset H_2(X, \mathbf{T})$. The canonical projection $H^2(X, \mathbf{T}) \rightarrow H^2(X, \mathbb{C})$ sends D_l to its class $[D_l]$ for $l = 1, \dots, m$. Since $\{[D_l]\}_{l=n+1}^m$ forms a basis of $H^2(X, \mathbb{C})$, we may choose a splitting $H^2(X, \mathbb{C}) \hookrightarrow H_{\mathbb{C}}^2(X, \mathbf{T})$ by taking the basic vector $[D_{n+k}]$ to D_{n+k} for $k = 1, \dots, m - n$. In this way we can regard $H^2(X, \mathbb{C})$ as a subspace of $H_{\mathbb{C}}^2(X, \mathbf{T})$.

Definition 3.2 (The mirror map). *Let X be a semi-Fano toric manifold. The mirror map is defined as the $1/z$ -coefficient of the I -function of X , which is an $H^2(X, \mathbb{C})$ -valued function in $\check{q} \in (\mathbb{C}^*)^{m-n}$.*

The $1/z$ -coefficient of $\sum_{d \in H_2^{\text{eff}}(X)} \check{q}^d I_d$ is of the form

$$-\sum_{l=1}^m g_l(\check{q}) D_l \in H_{\mathbb{C}}^2(X, \mathbf{T})$$

for some functions g_l in $\check{q} \in (\mathbb{C}^*)^{m-n}$, $l = 1, \dots, m$. Explicitly

$$g_l(\check{q}) := \sum_d \frac{(-1)^{(D_l \cdot d)} (-(D_l \cdot d) - 1)!}{\prod_{p \neq l} (D_p \cdot d)!} \check{q}^d$$

where the summation is over all curve classes $d \in H_2^{\text{eff}}(X)$ satisfying

$$-K_X \cdot d = 0, D_l \cdot d < 0 \text{ and } D_p \cdot d \geq 0 \text{ for all } p \neq l.$$

Then the mirror map is

$$\sum_{k=1}^{m-n} (\log \check{q}_k) [D_{n+k}] - \sum_{l=1}^m g_l(\check{q}) [D_l] = \sum_{k=1}^{m-n} (\log \check{q}_k - g^{\Psi_k}(\check{q})) [D_{n+k}]$$

where

$$(3.1) \quad g^{\Psi_k} := \sum_{l=1}^m (D_l, \Psi_k) g_l.$$

Thus in terms of the coordinates $(q_k)_{k=1}^{m-n}$ of $H^2(X, \mathbb{C})/2\pi\mathbf{i} H^2(X, \mathbb{Z})$, the mirror map $q(\check{q})$ is

$$(3.2) \quad q_k(\check{q}) = \check{q}_k \exp(-g^{\Psi_k}(\check{q}))$$

for $k = 1, \dots, m - n$.

While a priori $g_l(\check{q})$ is a formal power series in $\check{q}$ which can be considered as an element in the Novikov ring, by the theory of hypergeometric series it is known that $g_l(\check{q})$ is indeed convergent around $\check{q} = 0$ for each $l = 1, \dots, m$.

Moreover, the mirror map $q(\check{q})$ is a local diffeomorphism, and its inverse is denoted as $\check{q}(q)$.

Definition 3.3. The mirror complex moduli $\mathcal{M}^{\text{mir}}$ is defined as the domain of convergence of $(g_l(\check{q}))_{l=1}^m$ around $\check{q} = 0$ in $(\mathbb{C}^*)^{m-n}$.

The Kähler moduli is defined as the intersection of the complexified Kähler cone with the domain of convergence of the inverse mirror map $\check{q}(q)$. By abuse of notation we still denote the Kähler moduli as $\mathcal{K}_X^{\mathbb{C}}$.

The most important result in closed-string mirror symmetry for toric manifolds is the following:

Theorem 3.4 (Toric mirror theorem [18, 27]). *Let X be a compact semi-Fano toric manifold. Consider the J -function of X :*

$$(3.3) \quad J(q, z) = \exp \left(\frac{1}{z} \sum_{k=1}^{m-n} (\log q_k) [D_{n+k}] \right) \left(1 + \sum_a \sum_{d \in H_2^{\text{eff}}(X) \setminus \{0\}} q^d \left\langle 1, \frac{\phi_a}{z - \psi} \right\rangle_{0,2,d} \phi^a \right)$$

where $\{\phi_a\}$ is a homogeneous additive basis of $H^*(X)$ and $\{\phi^a\} \subset H^*(X)$ is its dual basis with respect to the Poincaré pairing. We always use $\langle \dots \rangle_{g,k,d}$ to denote the genus g , degree d descendent Gromov-Witten invariant of X with k insertions.

Then

$$I(\check{q}, z) = J(q(\check{q}), z)$$

where $q(\check{q})$ is the mirror map given in Definition 3.2.

In this paper, we are interested in open-string mirror symmetry. The Hori-Vafa superpotential (which plays a role analogous to that of the I -function in closed-string mirror symmetry) is the central object for this purpose.

Definition 3.5. *The Hori-Vafa superpotential of the toric manifold X is a holomorphic function $W^{\text{HV}} : \mathcal{M}^{\text{mir}} \times (\mathbb{C}^*)^n \rightarrow \mathbb{C}^*$ defined by*

$$W_{\check{q}}^{\text{HV}}(z_1, \dots, z_n) = z_1 + \dots + z_n + \sum_{k=1}^{m-n} \check{q}_k z^{v_{n+k}}$$

where $z^{v_{n+k}}$ denotes the monomial $\prod_{l=1}^n z_l^{(v_{n+k}, \nu_l)}$. It is pulled back to the Kähler moduli $\mathcal{K}_X^{\mathbb{C}}$ by substituting the inverse mirror map $\check{q}_k = \check{q}_k(q)$ ($k = 1, \dots, m-n$) in the above expression.

The Hori-Vafa potential may also be written as

$$(3.4) \quad \tilde{W}_{\check{q}}^{\text{HV}}(z_1, \dots, z_n) = \sum_{p=1}^n (\exp g_p(\check{q})) z_p + \sum_{k=1}^{m-n} (\exp g_{n+k}(\check{q})) \check{q}_k z^{v_{n+k}}.$$

via the coordinate change $z_p \mapsto (\exp g_p(\check{q})) z_p$, $p = 1, \dots, n$. Such a coordinate change will be necessary for the comparison with the disc potential (see Corollary 6.8).

The Hori-Vafa superpotential contains closed-string enumerative information of X :

Theorem 3.6 (Second form of the toric mirror theorem [18, 27]). *Let X be a semi-Fano toric manifold, $\omega \in \mathcal{K}_X^{\mathbb{C}}$ with coordinate $q(-\omega) = q$, and let W^{HV} be the Hori-Vafa superpotential. Then*

$$QH^*(X, \omega) \cong \text{Jac}(W_{\check{q}(q)}^{\text{HV}})$$

where $\check{q}(q)$ is the inverse mirror map. Moreover, the isomorphism is given by sending the generators $[D_{n+k}] \in QH^*(X, \omega)$ to $[\frac{\partial}{\partial \log q_k} W_{\check{q}(q)}^{\text{HV}}] \in \text{Jac}(W_{\check{q}(q)}^{\text{HV}})$.

3.2. Extended moduli. We have seen that I_d is indeed $H_{\mathbb{C}}^2(X, \mathbf{T})$ -valued. Thus it is natural to extend the mirror map and the Hori-Vafa potential $W_{\check{q}}^{\text{HV}}$ to $H_{\mathbb{C}}^2(X, \mathbf{T})$. (Extended moduli has been introduced by Iritani [22].)

Let

$$Q_l = \exp(-(\beta_l, \cdot)) \quad \text{for } l = 1, \dots, m$$

be the flat coordinates on $H_{\mathbb{C}}^2(X, \mathbf{T})/2\pi\mathbf{i} H^2(X, \mathbf{T})$. In terms of these coordinates, the canonical projection

$$H_{\mathbb{C}}^2(X, \mathbf{T})/2\pi\mathbf{i} H^2(X, \mathbf{T}) \rightarrow H^2(X, \mathbb{C})/2\pi\mathbf{i} H^2(X, \mathbb{Z})$$

is given by $q_k = Q^{-v_{n+k}} Q_{n+k}$ for $k = 1, \dots, m - n$. Here $Q^{-v_{n+k}}$ denotes the monomial $\prod_{l=1}^n Q_l^{(-v_{n+k}, \nu_l)}$. The splitting

$$H^2(X, \mathbb{C})/2\pi\mathbf{i} H^2(X, \mathbb{Z}) \hookrightarrow H_{\mathbb{C}}^2(X, \mathbf{T})/2\pi\mathbf{i} H^2(X, \mathbf{T})$$

is given by $Q_l = 1$ for $l = 1, \dots, n$ and $Q_l = q_l$ for $l = n + 1, \dots, m$.

Definition 3.7 (Extended moduli). *The extended Kähler moduli*

$$\tilde{\mathcal{K}}_X^{\mathbb{C}} \subset H_{\mathbb{C}}^2(X, \mathbf{T})/2\pi\mathbf{i} H^2(X, \mathbf{T})$$

is defined as the inverse image of the Kähler moduli $\mathcal{K}_X^{\mathbb{C}} \subset H^2(X, \mathbb{C})/2\pi\mathbf{i} H^2(X, \mathbb{Z})$ under the canonical projection

$$H_{\mathbb{C}}^2(X, \mathbf{T})/2\pi\mathbf{i} H^2(X, \mathbf{T}) \rightarrow H^2(X, \mathbb{C})/2\pi\mathbf{i} H^2(X, \mathbb{Z}).$$

The extended mirror complex moduli $\tilde{\mathcal{M}}^{\text{mir}}$ is the inverse image of $\mathcal{M}^{\text{mir}} \subset (\mathbb{C}^*)^{m-n}$ under the projection $(\mathbb{C}^*)^m \rightarrow (\mathbb{C}^*)^{m-n}$ defined by sending $(\check{Q}_1, \dots, \check{Q}_m) \in (\mathbb{C}^*)^m$ to $(\check{q}_1, \dots, \check{q}_{m-n}) \in (\mathbb{C}^*)^{m-n}$, $\check{q}_k = \check{Q}^{-v_{n+k}} \check{Q}_{n+k}$ for $k = 1, \dots, m - n$. $\mathcal{M}^{\text{mir}}$ can be regarded as a submanifold of $\tilde{\mathcal{M}}^{\text{mir}}$ by the splitting

$$\mathcal{M}^{\text{mir}} \hookrightarrow \tilde{\mathcal{M}}^{\text{mir}}$$

given by $\check{Q}_l = 1$ for $l = 1, \dots, n$ and $\check{Q}_{n+k} = \check{q}_k$ for $k = 1, \dots, m - n$.

Definition 3.8 (Extended superpotential and mirror map). *The extension of Hori-Vafa potential $W_{\check{q}}^{\text{HV}}$ from $\check{q} \in \mathcal{M}^{\text{mir}}$ to $\check{Q} \in \tilde{\mathcal{M}}^{\text{mir}}$ is defined as*

$$\mathcal{W}_{\check{Q}}^{\text{HV}}(z_1, \dots, z_n) := \sum_{l=1}^m \check{Q}_l z^{v_l}$$

where z^{v_l} denotes the monomial $\prod_{p=1}^n z_p^{(v_l, \nu_p)}$ (and so $z^{v_l} = z_l$ for $l = 1, \dots, n$).

The extended mirror map from $\tilde{\mathcal{M}}^{\text{mir}}$ to $\tilde{\mathcal{K}}_X^{\mathbb{C}}$ is defined to be

$$Q_l(\check{Q}) = \check{Q}_l \exp(-g_l(\check{q}(\check{Q})))$$

where $\check{q}(\check{Q})$ is the canonical projection $\check{q}_k = \check{Q}^{-v_{n+k}} \check{Q}_{n+k}$ for $k = 1, \dots, m - n$.

The extended inverse mirror map from $\tilde{\mathcal{K}}_X^{\mathbb{C}}$ to $\tilde{\mathcal{M}}^{\text{mir}}$ is defined to be

$$\check{Q}_l(Q) = Q_l \exp(g_l(\check{q}(q(Q))))$$

where $q(Q)$ is the canonical projection $q_k = Q^{-v_{n+k}} Q_{n+k}$ for $k = 1, \dots, m - n$, and $\check{q}(q)$ is the inverse mirror map.

There is a subtlety here: while $Q(\check{Q})$ and $\check{Q}(Q)$ are called the extended mirror map and extended inverse mirror map respectively, they are *not* necessarily inverse to each other! They are named in this way because they can be considered as extension of the mirror map and its inverse respectively in the following sense:

Proposition 3.9. *The extended mirror map is compatible with the mirror map. That is, the composition of the canonical projection $\tilde{\mathcal{K}}^{\mathbb{C}} \rightarrow \mathcal{K}^{\mathbb{C}}$ with the restriction of extended mirror map $Q(\check{Q})|_{\mathcal{M}^{\text{mir}}} : \mathcal{M}^{\text{mir}} \rightarrow \tilde{\mathcal{K}}^{\mathbb{C}}$ is equal to the mirror map $q(\check{q})$. (Recall that we always use the splitting in Definition 3.7 to regard $\mathcal{M}^{\text{mir}}$ as a submanifold of $\tilde{\mathcal{M}}^{\text{mir}}$.)*

Similarly, the extended inverse mirror map is compatible with the inverse mirror map in the sense that the composition of the canonical projection $\tilde{\mathcal{M}}^{\text{mir}} \rightarrow \mathcal{M}^{\text{mir}}$ with the restriction of extended inverse mirror map $\check{Q}(Q)|_{\mathcal{K}^{\mathbb{C}}} : \mathcal{K}^{\mathbb{C}} \rightarrow \tilde{\mathcal{M}}^{\text{mir}}$ is equal to the inverse mirror map $\check{q}(q)$.

As a consequence,

$$W_{\check{q}(q)}^{\text{HV}} = W_{\check{q}(\check{Q}(q))}^{\text{HV}} = \mathcal{W}_{\check{q}(\check{Q}(q))}^{\text{HV}}$$

where $\check{q}(q)$ is the inverse mirror map, $\check{Q}(q)$ is the restriction of the extended inverse mirror map $\check{Q}(Q)$ to the Kähler moduli $\mathcal{K}^{\mathbb{C}} \ni q$, and $\check{q}(\check{Q})$ is the canonical projection.

Proof. In terms of coordinates, the restriction of the extended mirror map to $\check{Q}_l = 1$ for $l = 1, \dots, n$, $\check{Q}_{n+k} = \check{q}_k$ for $k = 1, \dots, m - n$ is

$$\begin{aligned} Q_l &= \exp(-g_l(\check{q})) & \text{for } l = 1, \dots, n; \\ Q_{n+k} &= \check{q}_k \exp(-g_{n+k}(\check{q})) & \text{for } k = 1, \dots, m - n. \end{aligned}$$

Under the canonical projection $\tilde{\mathcal{K}}^{\mathbb{C}} \rightarrow \mathcal{K}^{\mathbb{C}}$,

$$\begin{aligned} q_k &= Q^{-v_{n+k}} Q_{n+k} \\ &= \check{q}_k \exp \left(- \left(g_{n+k}(\check{q}) + \sum_{p=1}^n (-v_{n+k}, \nu_p) g_p(\check{q}) \right) \right) \\ &= \check{q}_k \exp \left(- \left(\sum_{p=1}^m (\Psi_k, D_p) g_p(\check{q}_1, \dots, \check{q}_k) \right) \right) \end{aligned}$$

which is equal to the expression of the mirror map.

Similarly the restriction of the extended inverse mirror map to $Q_l = 1$ for $l = 1, \dots, n$, $Q_{n+k} = q_k$ for $k = 1, \dots, m - n$ is

$$\begin{aligned} \check{Q}_l &= \exp g_l(\check{q}(q)) & \text{for } l = 1, \dots, n; \\ \check{Q}_{n+k} &= q_k \exp g_{n+k}(\check{q}(q)) & \text{for } k = 1, \dots, m - n. \end{aligned}$$

Under the canonical projection $\tilde{\mathcal{M}}^{\text{mir}} \rightarrow \mathcal{M}^{\text{mir}}$,

$$\begin{aligned}\check{q}_k &= Q^{-v_{n+k}} Q_{n+k} \\ &= q_k \exp \left(g_{n+k}(\check{q}(q)) + \sum_{p=1}^n (-v_{n+k}, \nu_p) g_p(\check{q}(q)) \right) \\ &= q_k \exp \left(\sum_{p=1}^m (\Psi_k, D_p) g_p(\check{q}(q)) \right)\end{aligned}$$

which is equal to the expression of the inverse mirror map.

It is obvious that $W_{\check{q}}^{\text{HV}} = \mathcal{W}_{\check{Q}=\check{q}}^{\text{HV}}$ under the given splitting. Thus one has the equality relating the superpotential with its extended version given in this proposition. $\square$

Definition 3.10. An element $A = \sum_{l=1}^m \tilde{a}_l D_l \in H_{\mathbb{C}}^2(X, \mathbf{T})$ induces a differential operator

$$\hat{A}_{\check{Q}} := \sum_{l=1}^m \tilde{a}_l \partial_{\log \check{Q}_l}$$

which operates on functions on $\tilde{\mathcal{M}}^{\text{mir}}$. Such an association is linear, that is, $(c\hat{A} + \hat{B})_{\check{Q}} = c\hat{A}_{\check{Q}} + \hat{B}_{\check{Q}}$ for all $c \in \mathbb{C}$. The element A projects to $[A] \in H^2(X, \mathbb{C})$ which can be written as $\sum_{k=1}^{m-n} a_k [D_{n+k}]$ in terms of the basis $\{[D_{n+k}]\}_{k=1}^{m-n}$. It induces the differential operator

$$\hat{A}_{\check{q}} := \sum_{k=1}^{m-n} a_k \partial_{\log \check{q}_k}$$

which operates on functions on $\mathcal{M}^{\text{mir}}$. Similarly this association is linear.

Replacing $\check{Q}$ by Q and $\check{q}$ by q , A also induces the differential operators

$$\hat{A}_Q := \sum_{l=1}^m \tilde{a}_l \partial_{\log Q_l}$$

on the extended Kähler moduli $\tilde{\mathcal{K}}^{\mathbb{C}}$ and

$$\hat{A}_q := \sum_{k=1}^{m-n} a_k \partial_{\log q_k}$$

on the Kähler moduli $\mathcal{K}^{\mathbb{C}}$.

A good thing about the extension $\mathcal{W}^{\text{HV}}$ is the following observation:

Proposition 3.11. If $A, B \in H_{\mathbb{C}}^2(X, \mathbf{T})$ project to the same element in $H^2(X, \mathbb{C})$ (meaning that A and B are linearly equivalent), then

$$[\hat{A}_{\check{Q}} \mathcal{W}^{\text{HV}}] = [\hat{B}_{\check{Q}} \mathcal{W}^{\text{HV}}] \in \text{Jac}(\mathcal{W}_{\check{Q}}^{\text{HV}}).$$

In particular, restricting to the mirror moduli $\mathcal{M}^{\text{mir}}$, one has

$$[(\hat{A}_{\check{Q}} \mathcal{W}^{\text{HV}})(\check{q})] = [\hat{A}_{\check{q}} \mathcal{W}^{\text{HV}}] \in \text{Jac}(\mathcal{W}_{\check{q}}^{\text{HV}}).$$

Proof. For the first statement, it suffices to prove that if $\sum_{l=1}^m \tilde{a}_l D_l$ is linearly equivalent to zero, then

$$\sum_{l=1}^m \tilde{a}_l \partial_{\log \check{Q}_l} \mathcal{W}_{\check{Q}}^{\text{HV}}$$

is in the Jacobian ideal of $\mathcal{W}_{\check{Q}}^{\text{HV}}$.

Now

$$\sum_{l=1}^m \tilde{a}_l \partial_{\log \check{Q}_l} \mathcal{W}_{\check{Q}}^{\text{HV}} = \sum_{l=1}^m \tilde{a}_l \check{Q}_l z^{v_l}.$$

Since $\sum_{l=1}^m \tilde{a}_l D_l$ is linearly equivalent to zero, there exists $\nu \in M$ such that $(\nu, v_l) = \tilde{a}_l$ for all $l = 1, \dots, m$. Then the above expression is equal to

$$\sum_{l=1}^m (\nu, v_l) \check{Q}_l z^{v_l} = \frac{\partial}{\partial \log z^\nu} \mathcal{W}_{\check{Q}}^{\text{HV}}$$

where z^ν denotes the monomial $\prod_{l=1}^n z_l^{(\nu, v_l)}$.

For the second statement, write $A = \sum_{l=1}^m \tilde{a}_l D_l$ and its projection $[A] = \sum_{k=1}^{m-n} a_k [D_{n+k}]$. Then A and $\sum_{k=1}^{m-n} a_k D_{n+k} \in H_{\mathbb{C}}^2(X, \mathbf{T})$ projects to the same element in $H^2(X, \mathbb{C})$. Thus

$$[\hat{A}_{\check{Q}} \mathcal{W}^{\text{HV}}] = \left[\sum_{k=1}^{m-n} a_k \partial_{\log \check{Q}_{n+k}} \mathcal{W}^{\text{HV}} \right] \in \text{Jac}(\mathcal{W}_{\check{Q}}^{\text{HV}}).$$

Take $\check{Q}_l = 1$ for $l = 1, \dots, n$ and $\check{Q}_{n+k} = \check{q}_k$ for $k = 1, \dots, m-n$, since $\mathcal{W}_{\check{q}}^{\text{HV}} = W_{\check{q}}^{\text{HV}}$, we obtain

$$[(\hat{A}_{\check{Q}} \mathcal{W}^{\text{HV}})(\check{q})] = [\hat{A}_{\check{q}} W^{\text{HV}}] \in \text{Jac}(W_{\check{q}}^{\text{HV}}).$$

□

3.3. Batyrev elements. The toric mirror theorem 3.6 gives a presentation of the quantum cohomology ring. The generators of this presentation are given by the so called Batyrev elements defined as follows [3].

Definition 3.12 (Batyrev elements [3]). *The Batyrev elements, which are $H^2(X, \mathbb{C})$ -valued functions on $\mathcal{M}^{\text{mir}}$, are defined as follows. For $k = 1, \dots, m-n$,*

$$B_{n+k} := \sum_{r=1}^{m-n} \frac{\partial \log q_r(\check{q})}{\partial \log \check{q}_k} [D_{n+r}]$$

where $q(\check{q})$ is the mirror map. For $l = 1, \dots, n$,

$$B_l := \sum_{k=1}^{m-n} (B_l, \Psi_k) B_{n+k}.$$

A key step to prove the toric mirror theorem 3.6 is to show that the Batyrev elements satisfy two sets of explicit relations [3]:

- (1) Linear relations. It follows from the definition that $\{B_l\}_{l=1}^m$ satisfies the same linear relations as that satisfied by $\{D_l\}_{l=1}^m$, namely, for every $\nu \in M$,

$$\sum_{l=1}^m (\nu, v_l) B_l = 0.$$

- (2) Multiplicative relations. For every $k = 1, \dots, m - n$,

$$B_1^{(\Psi_k, D_1)} * \dots * B_m^{(\Psi_k, D_m)} = q_k$$

where $B_l^{(\Psi_k, D_l)}$ means B_l quantum-multiplies itself for (Ψ_k, D_l) times. This relation is the essential part of the toric mirror theorem (second form, see Theorem 3.6).

Batyrev elements can also be lifted to $\tilde{\mathcal{M}}^{\text{mir}}$:

Definition 3.13 (Extended Batyrev elements). *For $l = 1, \dots, m$, define*

$$\tilde{B}_l := \sum_{p=1}^m \frac{\partial \log Q_p(\check{Q})}{\partial \log \check{Q}_l} D_p$$

which is an $H_{\mathbb{C}}^2(X, \mathbf{T})$ -valued function on $\tilde{\mathcal{M}}^{\text{mir}}$. Here $Q(\check{Q})$ is the extended mirror map given in Definition 3.8.

Proposition 3.14. *For $l = 1, \dots, m$, we have*

$$[\tilde{B}_l|_{\mathcal{M}^{\text{mir}}}] = B_l.$$

Proof. In terms of the basis $\{[D_{n+1}], \dots, [D_m]\}$ of $H^2(X, \mathbb{C})$, $[D_p] = \sum_{r=1}^{m-n} (D_p, \Psi_r) [D_{n+r}]$. Hence

$$\begin{aligned} [\tilde{B}_l] &= \sum_{p=1}^m \frac{\partial \log Q_p(\check{Q})}{\partial \log \check{Q}_l} \sum_{r=1}^{m-n} (D_p, \Psi_r) [D_{n+r}] \\ &= \sum_{r=1}^{m-n} \frac{\partial \log q_r(Q(\check{Q}))}{\partial \log \check{Q}_l} [D_{n+r}] \end{aligned}$$

where $q_r(Q) = Q^{-v_r} Q_{n+k}$ for $r = 1, \dots, m - n$ is the canonical projection. Recall that the extended mirror map is $\log Q_p(\check{Q}) = \log \check{Q}_p - g_p(\check{q}(\check{Q}))$ where $\check{q}(\check{Q})$ is the canonical projection

given by $\check{q}_r(\check{Q}) = \check{Q}^{-v_r} \check{Q}_{n+k}$ for $r = 1, \dots, m - n$. Then

$$\begin{aligned}
[\tilde{B}_l] &= \sum_{r=1}^{m-n} \frac{\partial \log q_r(Q(\check{Q}))}{\partial \log \check{Q}_l} [D_{n+r}] \\
&= \sum_{r=1}^{m-n} \sum_{p=1}^m \frac{\partial \log q_r}{\partial \log Q_p} \frac{\partial \log Q_p}{\partial \log \check{Q}_l} [D_{n+r}] \\
&= \sum_{r=1}^{m-n} \sum_{p=1}^m (D_p, \Psi_r) \left(\delta_{pl} - \frac{\partial g_p(\check{q}(\check{Q}))}{\partial \log \check{Q}_l} \right) [D_{n+r}] \\
&= \sum_{r=1}^{m-n} \left((D_l, \Psi_r) - \frac{\partial g^{\Psi_r}(\check{q}(\check{Q}))}{\partial \log \check{Q}_l} \right) [D_{n+r}] \\
&= \sum_{r=1}^{m-n} \left((D_l, \Psi_r) - \sum_{k=1}^{m-n} \frac{\partial g^{\Psi_r}}{\partial \log \check{q}_k} \frac{\partial \log \check{q}_k}{\partial \log \check{Q}_l} \right) [D_{n+r}] \\
&= \sum_{r=1}^{m-n} \left((D_l, \Psi_r) - \sum_{k=1}^{m-n} (\Psi_k, D_l) \frac{\partial g^{\Psi_r}}{\partial \log \check{q}_k}(\check{q}(\check{Q})) \right) [D_{n+r}]
\end{aligned}$$

where $g^{\Psi_r} = \sum_{p=1}^m (D_p, \Psi_r) g_p$. Now restricting $\check{Q}_1 = \dots = \check{Q}_n = 1$ and $\check{Q}_{n+k} = \check{q}_{n+k}$ for $k = 1, \dots, m - n$, and since the extended mirror map is compatible with the mirror map (see Proposition 3.9), the above equals

$$\sum_{r=1}^{m-n} \sum_{k=1}^{m-n} (\Psi_k, D_l) \frac{\partial \log q_r}{\partial \log \check{q}_k}(\check{q}) [D_{n+r}] = B_l.$$

□

It follows from the above discussions that Batyrev elements have an extremely simple form under the isomorphism $QH^*(X, \omega_q) \cong \text{Jac}(W_{\check{q}(q)}^{\text{HV}})$:

Proposition 3.15. *Under the isomorphism $QH^*(X, \omega_q) \cong \text{Jac}(W_{\check{q}(q)}^{\text{HV}})$ of Theorem 3.6, the Batyrev elements B_l are mapped to z^{v_l} for $l = 1, \dots, n$, and $\check{q}_{l-n}(q) z^{v_l}$ for $l = n + 1, \dots, m$.*

Equivalently, each B_l is mapped to $(\exp g_l(\check{q}(q))) Z_l$ for $l = 1, \dots, m$ under $QH^*(X, \omega_q) \cong \text{Jac}(W_{\check{q}(q)}^{\text{HV}})$, where Z_l is defined by Equation (2.2).

Proof. Associate $D_p \in H_{\mathbb{C}}^2(X, \mathbf{T})$ to the differential operator $\frac{\partial}{\partial \log Q_p}$ for $p = 1, \dots, m$. Then $\tilde{B}_l$ is associated with

$$\widehat{\tilde{B}_l} = \sum_{p=1}^m \frac{\partial \log Q_p}{\partial \log \check{Q}_l} \frac{\partial}{\partial \log Q_p} = \frac{\partial}{\partial \log \check{Q}_l}.$$

Restricting to $\mathcal{M}^{\text{mir}}$, by Proposition 3.14 $\tilde{B}_l$ projects to $B_l \in H^2(X, \mathbb{C})$. By Proposition 3.11,

$$[\widehat{\tilde{B}_l} W^{\text{HV}}(\check{q}(q))] = [\widehat{B}_l W_{\check{q}(q)}^{\text{HV}}] \in \text{Jac}(W_{\check{q}(q)}^{\text{HV}}).$$

On the other hand,

$$\widehat{B}_l \mathcal{W}^{\text{HV}}(\check{q}(q)) = \frac{\partial}{\partial \log \check{Q}_l} \sum_{p=1}^m \check{Q}_p z^{v_p} = \check{Q}_l z^{v_l}.$$

Thus

$$[\widehat{B}_l \mathcal{W}_{\check{q}(q)}^{\text{HV}}] = z^{v_l}$$

for $l = 1, \dots, n$ and

$$[\widehat{B}_{n+k} \mathcal{W}_{\check{q}(q)}^{\text{HV}}] = \check{q}_k(q) z^{v_{n+k}}$$

for $k = 1, \dots, m - n$.

Under the coordinate change (3.4), z_l changes to $(\exp g_l(\check{q}(q)))z_l$ for $l = 1, \dots, n$, and for $l = n + 1, \dots, m$, $\check{q}_{l-n}(q)z^{v_l}$ changes to

$$\begin{aligned} \check{q}_{l-n}(q) \prod_{p=1}^n ((\exp g_p(\check{q}(q)))z_p)^{(v_l, \nu_p)} &= z^{v_l} q_{l-n}(\exp g^{\Psi_{l-n}}(\check{q}(q))) \prod_{p=1}^n (\exp g_p(\check{q}(q))) \\ &= q_{l-n}(\exp g_l(\check{q}(q)))z^{v_l}. \end{aligned}$$

□

4. SEIDEL REPRESENTATIONS FOR TORIC MANIFOLDS

In this Section we review the construction and properties of Seidel representations in the toric case. A key insight of this paper is that open Gromov-Witten invariants of a toric manifold X are equal to some closed Gromov-Witten invariants of certain manifolds related to X . These manifolds are used to construct the so-called Seidel representation [32, 30], which is an action⁵ of $\pi_1(\text{Ham}(X, \omega))$ on $QH^*(X, \omega)$, and so we call them the Seidel spaces in this paper.

Definition 4.1. *Let X be a manifold. Suppose that we have an action $\rho : \mathbb{C}^* \times X \rightarrow X$ of $\mathbb{C}^*$ on X . The manifold*

$$E = E_\rho := (X \times (\mathbb{C}^2 \setminus \{0\})) / \mathbb{C}^*$$

is called the Seidel space associated to the action ρ , where $z \in \mathbb{C}^*$ acts on the second factor $\mathbb{C}^2 \setminus \{0\} \ni (u, v)$ by $z \cdot (u, v) = (zu, zv)$.

The Seidel space E is an X -bundle over $\mathbb{P}^1$ where the bundle map $(X \times (\mathbb{C}^2 \setminus \{0\})) / \mathbb{C}^* \rightarrow \mathbb{P}^1$ is given by the projection to the second factor.

Let X be a toric manifold of complex dimension n , given by a fan Σ^X supported in the vector space $N \otimes_{\mathbb{Z}} \mathbb{R}$, where N is a lattice. Each lattice point $v \in N$ gives a $\mathbb{C}^*$ -action on X . In particular, the minimal generator $v_j \in N$ of a ray of the fan Σ^X gives a $\mathbb{C}^*$ -action and thus defines a corresponding Seidel space $E = E_j$. It is a toric manifold of dimension $n + 1$ whose fan Σ^E has rays generated by $v_l^E = (0, v_l)$ for $l = 1, \dots, m$, $v_0^E = (1, 0)$ and $v_\infty^E = (-1, v_j)$.

On the other hand, we may also use the opposite direction $-v_j \in N$ to generate a $\mathbb{C}^*$ -action. The corresponding Seidel space will be denoted by $E^- = E_j^-$. It is also a toric manifold of dimension $n + 1$ whose fan Σ^{E^-} has rays generated by $v_l^{E^-} = (0, v_l)$ for $l = 1, \dots, m$, $v_0^{E^-} = (1, 0)$ and $v_\infty^{E^-} = (-1, -v_j)$.

⁵ $\text{Ham}(X, \omega)$ denotes the group of Hamiltonian diffeomorphisms on (X, ω) .

One of the main results in this paper is that, the open invariants of X are closely related to the open invariants of the Seidel spaces $E^- = E_j^-$, $j = 1, \dots, m$. Since E^- is a toric manifold, $\pi_2(E^-, \mathbf{T}^{E^-})$ is generated by the basic disc classes which are denoted as b_l , $l = 0, 1, \dots, m, \infty$ (while recall that basic disc classes in X are denoted as β_l for $l = 1, \dots, m$). Moreover, the toric prime divisors of E^- are denoted as $\mathcal{D}_0, \mathcal{D}_1, \dots, \mathcal{D}_m, \mathcal{D}_\infty$ (while recall that toric prime divisors of X are denoted as D_l for $l = 1, \dots, m$).

Viewing Seidel spaces as X -bundles over $\mathbb{P}^1$, one has the following specific sections of the Seidel spaces:

Definition 4.2. *Let X be a toric manifold, and let $v_j \in N$ be the minimal generator of a ray in the fan of X . Let $E = E_j$ and $E^- = E_j^-$ be the Seidel spaces associated to v_j and $-v_j$ respectively. Under the $\mathbb{C}^*$ -action generated by either v_j or $-v_j$, there are two fixed loci in X . One of them is $D_j \subset X$, and each point $p \in D_j$ gives a section $\sigma = \sigma_j : \mathbb{P}^1 \rightarrow E$ ($\sigma^- = \sigma_j^- : \mathbb{P}^1 \rightarrow E^-$ resp.) whose value is constantly $p \in D_j \subset X$. It is called the zero section of E (E^- resp.). Similarly each point p in another fixed locus S gives a section $\sigma_\infty : \mathbb{P}^1 \rightarrow E$ ($\sigma_\infty^- : \mathbb{P}^1 \rightarrow E^-$ resp.) whose value is constantly $p \in S \subset X$, and it is called a infinity section of E (E^- resp.).*

The various notations in the above definition are illustrated by the following figure of the Seidel spaces of $\mathbb{P}^1$.




FIGURE 2. This figure shows the Seidel spaces E and E^- associated to the two different torus actions on $\mathbb{P}^1$ and their sections corresponding to the fixed points in $\mathbb{P}^1$ under the action. In this simple case both E and E^- are the Hirzebruch surface $\mathbf{F}_1$, while in general they are different manifolds.

By abuse of notation their classes in $H_2(E)$ and $H_2(E^-)$ are also denoted as σ , σ_∞ and σ^- , σ_∞^- respectively. Notice that under the $\mathbb{C}^*$ -action generated by v_j , D_j has negative weight (namely, -1); while under the $\mathbb{C}^*$ -action generated by $-v_j$, the other fixed loci has negative weight. Then using Lemma 2.2 of González-Iritani [20], all the curve classes of E (E^- resp.) are generated by σ (σ_∞^- resp.) and curve classes of X :

Proposition 4.3 (Lemma 2.2 of [20]).

$$H_2^{\text{eff}}(E) = \mathbb{Z}_{\geq 0}[\sigma] + H_2^{\text{eff}}(X)$$

where $H_2^{\text{eff}}(E)$ and $H_2^{\text{eff}}(X)$ denote the Mori cones of effective curve classes of E and X respectively. Similarly,

$$H_2^{\text{eff}}(E^-) = \mathbb{Z}_{\geq 0}[\sigma_\infty^-] + H_2^{\text{eff}}(X)$$

where $H_2^{\text{eff}}(E)$ and $H_2^{\text{eff}}(X)$ denote the Mori cones of effective curve classes of E and X respectively.

The section class $\sigma^- \in H_2^{\text{eff}}(E^-)$ can also be written as $\sigma^- = b_0 + b_\infty + b_j$, where we recall that b_l for $l = 0, 1, \dots, m, \infty$ are the basic disc classes of E^- . It is the most important curve class to us as we shall see in the next section. By Proposition 4.3, it can also be written as

$$\sigma^- = \sigma_\infty^- + \underline{C}$$

for some curve class $\underline{C}$ in X .

The Seidel element corresponding to a $\mathbb{C}^*$ -action on X is defined as follows.

Definition 4.4 (Seidel element). *Given a $\mathbb{C}^*$ -action on X , let E be the corresponding Seidel space. The Seidel element is*

$$S := \sum_a \sum_{d \in H_2^{\text{sec}}(E)} q^d \langle \phi_a^E \rangle_{0,1,d} \phi^a = q^\sigma \sum_a \sum_{\alpha \in H_2^{\text{eff}}(X)} q^\alpha \langle \phi_a^E \rangle_{0,1,\sigma+\alpha} \phi^a$$

where $\{\phi_a\}$ is a basis of $H^*(X)$ and $\{\phi^a\}$ is the dual basis with respect to Poincaré pairing. $\phi_a^E \in H^*(E)$ denotes the push-forward of ϕ_a under the inclusion of X into E as a fiber. Moreover,

$$H_2^{\text{sec}}(E) := \{\sigma + \alpha : \alpha \in H_2^{\text{eff}}(X)\} \subset H_2^{\text{eff}}(E)$$

where σ is the section class of E corresponding to the fixed locus in X with negative weight.

The normalized Seidel element is

$$S^\circ := \sum_a \sum_{\alpha \in H_2^{\text{eff}}(X)} q^\alpha \langle \phi_a^E \rangle_{0,1,\sigma+\alpha} \phi^a \in QH^*(X, \omega_q).$$

Using degeneration arguments, Seidel [32] (in monotone case) and McDuff [30] proved that if ρ_1, ρ_2 are two commuting $\mathbb{C}^*$ -actions and $\rho_3 = \rho_1 * \rho_2$ is the composition, then the corresponding Seidel elements S_1, S_2, S_3 satisfy the relation⁶

$$S_3 = S_1 * S_2,$$

(here $*$ denotes the quantum multiplication) under the following assignment of relations between Novikov variables of E_1, E_2, E_3 .

The $\mathbb{C}^*$ -actions ρ_1, ρ_2 define an X -bundle $\hat{E}$ over $\mathbb{P}^1 \times \mathbb{P}^1$:

$$\hat{E} := (X \times (\mathbb{C}^2 \setminus \{0\}) \times (\mathbb{C}^2 \setminus \{0\})) / (\mathbb{C}^* \times \mathbb{C}^*)$$

where $(z_1, z_2) \in \mathbb{C}^* \times \mathbb{C}^*$ acts by

$$(z_1, z_2) \cdot (x, (u_1, v_1), (u_2, v_2)) := (\rho_1(z_1, \rho_2(z_2, x)), (z_1 u_1, z_1 v_1), (z_2 u_2, z_2 v_2)).$$

$\hat{E}$ restricted to $\mathbb{P}^1 \times \{[0, 1]\}$ is the Seidel space E_1 associated to ρ_1 ; $\hat{E}$ restricted to $\{[1, 0]\} \times \mathbb{P}^1$ is the Seidel space E_2 associated to ρ_2 ; $\hat{E}$ restricted to the diagonal $\{(p_1, p_2) \in \mathbb{P}^1 \times \mathbb{P}^1 : p_1 = p_2\}$ is the Seidel space E_3 associated to the composition ρ_3 . Effective curves classes in $\hat{E}$ are generated by those of E_1 and E_2 . In particular, σ_3 , the section class of E_3 which corresponds to the fixed locus in X with negative weight under the action of ρ_3 , when pushed forward to

⁶Indeed they proved this in a more general situation in which the Seidel elements are generated by loops in $\pi_1(\text{Ham}(X, \omega))$. Then every element in $\pi_1(\text{Ham}(X, \omega))$ gives a Seidel element which acts on $QH^*(X)$ by quantum multiplication, and they showed that it defines an action of $\pi_1(\text{Ham}(X, \omega))$ on $QH^*(X)$.

a curve class in $\hat{E}$, is of the form $d_1 + d_2$ for some $d_1 \in H_2^{\text{eff}}(E_1)$ and $d_2 \in H_2^{\text{eff}}(E_2)$. Then assign the relation

$$q^{\sigma_3} := q^{d_1} q^{d_2}$$

between the Novikov variables q^{d_1} , q^{d_2} , q^{σ_3} of E_1, E_2 and E_3 respectively. The Novikov variables q^α of E_i , where $\alpha \in H_2^{\text{eff}}(X)$, are set to equal for $i = 1, 2, 3$. Thus once the Novikov variables of E_1 and E_2 are fixed (with the requirement that q^α are the same for all $\alpha \in H_2^{\text{eff}}(X)$), those of E_3 are automatically fixed.

Now coming back to our case that ρ_1 is the action generated by v_j and ρ_2 is the action generated by $-v_j$, their composition is the trivial action. We write S_j and S_j^- for the two Seidel elements. The Seidel element generated by the trivial action is simply q^{σ_0} , where σ_0 is a section of the trivial bundle $X \times \mathbb{P}^1 \rightarrow \mathbb{P}^1$. The above assignment of Novikov variables gives

$$q^{\sigma_0} = q^{\sigma_j + \sigma_j^-}.$$

Hence one has the following equality for Seidel representation:

Proposition 4.5.

$$S_j * S_j^- = q^{\sigma_j + \sigma_j^-}.$$

5. RELATING OPEN AND CLOSED INVARIANTS

Open Gromov-Witten invariants are difficult to compute in general, because there are highly nontrivial obstructions to the moduli problems, and in contrast to closed Gromov-Witten theory, localization techniques cannot be applied. In [4, 23], under certain restrictions on the geometry of the toric manifold X , it was shown that open Gromov-Witten invariants appearing in the superpotential of X were equal to certain closed Gromov-Witten invariants of X (or certain compactification of X when X is non-compact). This gives an effective way to compute open Gromov-Witten invariants because closed Gromov-Witten invariants can be computed by various techniques such as localization.

However, for an arbitrary toric manifold X , the geometric technique in [4, 23] fails, and searching for spaces whose closed Gromov-Witten invariants correspond to open Gromov-Witten invariants of X becomes more difficult. An exciting discovery in this paper is that Seidel spaces associated to X , which are one dimensional higher than X , are indeed what we need in order to have such an open-closed comparison. Moreover it works for *all* semi-Fano toric manifolds. Namely,

Theorem 5.1. *Let X be a semi-Fano toric manifold and $\beta \in \pi_2(X, \mathbf{T}^X)$ a disc class of Maslov index two bounded by a Lagrangian torus fiber $\mathbf{T}^X \subset X$. Then β must be of the form $\beta_j + \alpha$ for some basic disc class β_j of X ($j=1, \dots, m$) and $\alpha \in H_2^{\text{eff}}(X)$ with $c_1(\alpha) = 0$.*

Let $v_j \in N$ be the minimal generator of the ray in the fan of X corresponding to the basic disc class β_j . Let $E^- = E_j^-$ be the Seidel space corresponding to the $\mathbb{C}^*$ -action generated by $-v_j = -\partial\beta_j$, and denote by $\mathbf{T}^{E^-}$ a Lagrangian torus fiber of E^- . Any class $a \in H^*(X)$ can be pushed forward (via Poincaré duality) by the inclusion $X \hookrightarrow E^-$ of X as a fiber to give a class in $H^*(E^-)$, and it is denoted as a^{E^-} . Then for every toric prime divisor $D = D_i$ of X , one has

$$n_{1,1}^X(\beta; D; [\text{pt}]_{\mathbf{T}^X}) = \langle D^{E^-}, [\text{pt}]_{E^-} \rangle_{0,2,\sigma^- + \alpha^{E^-}}^{E^-}$$

where $\sigma^- \in H_2(E^-)$ is the zero section class of E^- (see Definition 4.2), $[\text{pt}]_{\mathbf{T}^X} \in H^n(\mathbf{T}^X)$ is a point class of $\mathbf{T}^X$ and $[\text{pt}]_{E^-} \in H^{2n}(E^-)$ is a point class of E^- .

The statement that a stable disc class of Maslov index two bounded by $\mathbf{T}^X$ is of the form $\beta_j + \alpha$ was proved by Cho-Oh [10] and Fukaya-Oh-Ohta-Ono [14], and it is recalled in Lemma 2.3. We also need the following simple lemma about curves in the Seidel space E^- representing α :

Lemma 5.2. *Assume the setting as in Theorem 5.1. Let $C \subset E^-$ be a rational curve representing a fiber class of $E^- \rightarrow \mathbb{P}^1$ with $c_1(C) \leq 1$. Then C is contained in the union of the toric divisors of E^- , i.e. $C \subset \bigcup_{l=1}^m \mathcal{D}_l$.*

Proof. Since C represents a fiber class (i.e. a class in $H_2^{\text{eff}}(X)$), its image under $E^- \rightarrow \mathbb{P}^1$ can only be a point, which means C belongs to a fiber of $E^- \rightarrow \mathbb{P}^1$, which is identified as X . Note that the Chern number of C in X is the same as that of C in E^- . Since X is semi-Fano, every component of C has non-negative Chern number. But $c_1(C) \leq 1$. Thus each component of C has $c_1 \leq 1$. Let $C' \subset C$ be a component. Then $-K_X \cdot C'$ is either 0 or 1. Suppose $-K_X \cdot C' = 0$. It is impossible to have $D_i \cdot C' = 0$ for all i since this means $[C'] = 0$. So there exists an i such that $D_i \cdot C' < 0$, which implies that $C' \subset D_i$. Suppose $-K_X \cdot C' = 1$. It is possible that $D_i \cdot C' < 0$ for some i , which implies $C' \subset D_i$. The other possibility is that $D_i \cdot C' = 1$ for some i and $D_j \cdot C' = 0$ for $j \neq i$. This is impossible since it violates linear relations. We thus conclude that every component of C lies in a toric divisor of X . Under the inclusion $X \hookrightarrow E^-$ as a fiber, $D_l \subset \mathcal{D}_l$ for all $l = 1, \dots, m$. Thus $C \subset \bigcup_{l=1}^m \mathcal{D}_l$. $\square$

The rest of this section is devoted to prove Theorem 5.1. The proof is divided into two main steps. First, we equate the open invariant $n_{1,1}(\beta; D; [\text{pt}]_{\mathbf{T}^X})$ of X to certain open invariant of $E^- = E_j^-$ (Theorem 5.3). Then we show that this open invariant of E^- is equal to the closed invariant $\langle D^{E^-}, [\text{pt}]_{\mathbf{T}^{E^-}} \rangle_{0,2,\sigma^-+\alpha}$ of E^- (Theorem 5.6). Here $D^{E^-} \in H^4(E^-)$ is the pushforward of $D \in H^2(X)$ under the inclusion $X \hookrightarrow E^-$ of X as a fiber. Since D is a divisor of X , D^{E^-} is of complex codimension two in E^- .

5.1. First step. The precise statement of the first main step is the following:

Theorem 5.3. *Assume the notations as in Theorem 5.1. Then*

$$n_{1,1}^X(\beta_j + \alpha; D; [\text{pt}]_{\mathbf{T}^X}) = n_{1,1}^{E^-}(b_0 + b_j + \alpha; D^{E^-}; [\text{pt}]_{\mathbf{T}^{E^-}})$$

where we recall that b_l ($l = 0, 1, \dots, m, \infty$) are the basic disc classes of E^- (see Section 4). Moreover $[\text{pt}]_{\mathbf{T}^{E^-}} \in H^{n+1}(\mathbf{T}^{E^-})$ denotes the point class of the Lagrangian torus fiber of E^- .

Recall that

$$n_{1,1}^X(\beta_j + \alpha; D; [\text{pt}]_{\mathbf{T}^X}) := ([\mathcal{M}_{1,1}^{\text{op}}(\beta_j + \alpha; D)]_{\text{virt}}, [\text{pt}]_{\mathbf{T}^X}) \in \mathbb{Q}$$

and by definition of Poincaré pairing, this is the same as

$$\iota_{\mathbf{T}^X}^*[\mathcal{M}_{1,1}^{\text{op}}(\beta_j + \alpha; D)]_{\text{virt}} \in H^0(\text{pt}, \mathbb{Q}) \cong \mathbb{Q}.$$

where $\iota_{\mathbf{T}^X} : \{\text{pt}\} \hookrightarrow \mathbf{T}^X$ is an inclusion of a point to $\mathbf{T}^X$. Similarly

$$n_{1,1}^{E^-}(b_0 + b_j + \alpha; D^{E^-}; [\text{pt}]_{\mathbf{T}^{E^-}}) = \iota_{\mathbf{T}^{E^-}}^*[\mathcal{M}_{1,1}^{\text{op}}(b_0 + b_j + \alpha; D^{E^-})]_{\text{virt}} \in H^0(\text{pt}, \mathbb{Q}) \cong \mathbb{Q}$$

where $\iota_{\mathbf{T}^{E^-}} : \{\text{pt}\} \hookrightarrow \mathbf{T}^{E^-}$ is an inclusion of a point to $\mathbf{T}^{E^-}$. We denote the images of $\iota_{\mathbf{T}^X}$ and $\iota_{\mathbf{T}^{E^-}}$ to be $\text{pt}_{\mathbf{T}^X}$ and $\text{pt}_{\mathbf{T}^{E^-}}$ respectively.

By Lemma 2.3, a stable disc in $\mathcal{M}_{1,1}^{\text{op}}(\beta; D)$ has only one disc component. Thus it never splits into the union of two stable discs. Hence $\mathcal{M}_{1,1}^{\text{op}}(\beta; D)$ has no codimension-one boundary. The following lemma shows that $\mathcal{M}_{1,1}^{\text{op}}(b_0 + b_j + \alpha; D^{E^-})$ also has this property, whose proof requires a more careful analysis of the stable discs since $b_0 + b_j$ has Maslov index four (which is not the minimal Maslov index of $\mathbf{T}^{E^-}$), and E^- might not be semi-Fano:

Lemma 5.4. *Assume the settings as above. A stable disc in $\mathcal{M}_{1,1}^{\text{op}}(b_0 + b_j + \alpha; D^{E^-})$ consists of a holomorphic disc component and a rational curve, which are attached to each other at only one nodal point. The disc component belongs to the class $b_0 + b_j$ for some $j = 1, \dots, m$, and the rational curve belongs to α . In particular, $\mathcal{M}_{1,1}^{\text{op}}(b_0 + b_j + \alpha; D^{E^-})$ has no codimension-one boundary.*

Proof. Consider a stable disc ϕ in $\mathcal{M}_{1,1}^{\text{op}}(b_0 + b_j + \alpha; D^{E^-})$. It consists of several disc components and sphere components. Notice that $(b_0 + b_j + \alpha, \mathcal{D}_\infty) = 0$, where $(\cdot, \cdot)$ denotes the Poincaré pairing on $H^*(E^-)$. Since every holomorphic disc bounded by $\mathbf{T}^{E^-}$ and every holomorphic sphere in E^- has non-negative intersection with $\mathcal{D}_\infty$, this implies that each sphere component of ϕ has intersection number 0 with $\mathcal{D}_\infty$. So every sphere component of ϕ is in a fiber class as otherwise it would have positive intersection number with $\mathcal{D}_\infty$. In particular each sphere component of ϕ has non-negative Chern number and is contained in a fiber of $E^- \rightarrow \mathbb{P}^1$. Together with the fact that ϕ has Maslov index $\mu(b_0) + \mu(b_j) + 2c_1(\alpha) = 4$, this implies that each disc component has at most Maslov index four.

Suppose a disc component of ϕ has Maslov index four. Then all the sphere components have Chern number zero. Since every non-constant holomorphic disc has Maslov index at least two, the other disc components of ϕ must be constant, and they are mapped to $\mathbf{T}^{E^-}$. On the other hand the interior marked point p^{int} of ϕ has to be mapped to D^{E^-} , which is disjoint from $\mathbf{T}^{E^-}$. Hence p^{int} cannot be located in the constant disc components. But then at least one of the constant disc component does not have three special points, and hence making ϕ unstable. This shows that ϕ has only one disc component which has Maslov index four.

Then we prove that the disc component is attached with the holomorphic spheres at only one interior nodal point. Holomorphic discs bounded by a Lagrangian torus fiber have been classified by Cho-Oh [10]. In particular if a holomorphic disc of Maslov index four passes through D^{E^-} , it intersects with the union of toric divisors at only one single interior point. On the other hand, by Lemma 5.2, all the sphere components are mapped to the union of the toric divisors $\mathcal{D}_l$, $l = 1, \dots, m$. Thus the disc component is attached with the holomorphic spheres at only one interior nodal point. This implies that ϕ is the union of a holomorphic disc and a rational curve joint at a single nodal point. Since the holomorphic disc represents $b_0 + b_j$, the rational curve must represent α .

Now suppose otherwise that every disc component of ϕ has Maslov index less than four. Then ϕ must have a disc component of Maslov index two. Then the other disc components have Maslov index at most two, and the sphere components have Chern number at most one.

Moreover the sphere components belong to some fiber classes. By Lemma 5.2, each of them is contained in $\mathcal{D}_l$ for some $l = 1, \dots, m$.

A holomorphic disc of Maslov index at most two does not pass through D^{E^-} . Thus the interior marked point p^{int} of ϕ must be located in a sphere component. But $\phi(p^{\text{int}}) \in D^{E^-}$ which is contained in the fiber at 0. This implies that this sphere component is contained in $\mathcal{D}_0 \cap \mathcal{D}_l$ for some $l = 1, \dots, m$. However, a holomorphic disc of Maslov index at most two does not pass through $\mathcal{D}_0 \cap \mathcal{D}_l$, and so none of the disc components is connected to this sphere component. We thus conclude that this situation cannot occur.

We have now proved that ϕ has only one disc component. This implies that it never splits into two stable discs, meaning disc bubbling never occurs. Thus the moduli $\mathcal{M}_{1,1}^{\text{op}}(b_0 + b_j + \alpha; D^{E^-})$ has no codimension-one boundary. $\square$

Now both $\mathcal{M}_{1,1}^{\text{op}}(\beta_j + \alpha; D)$ and $\mathcal{M}_{1,1}^{\text{op}}(b_0 + b_j + \alpha; D^{E^-})$ have no codimension-one boundaries. By [12, Lemma A1.43], we have

$$\begin{aligned} n_{1,1}^X(\beta_j + \alpha; D; [\text{pt}]_{\mathbf{T}^X}) &= \iota_{\mathbf{T}^X}^*[\mathcal{M}_{1,1}^{\text{op}}(\beta_j + \alpha; D)]_{\text{virt}} \\ &= [\mathcal{M}_{1,1}^{\text{op}}(\beta_j + \alpha; D; \text{pt}_{\mathbf{T}^X})]_{\text{virt}} \in H^0(D \times \{\text{pt}_{\mathbf{T}^X}\}, \mathbb{Q}) = \mathbb{Q} \end{aligned}$$

and

$$\begin{aligned} (5.1) \quad n_{1,1}^{E^-}(b_0 + b_j + \alpha; D^{E^-}; [\text{pt}]_{\mathbf{T}^{E^-}}) &= \iota_{\mathbf{T}^{E^-}}^*[\mathcal{M}_{1,1}^{\text{op}}(b_0 + b_j + \alpha; D^{E^-})]_{\text{virt}} \\ &= [\mathcal{M}_{1,1}^{\text{op}}(b_0 + b_j + \alpha; D^{E^-}; \text{pt}_{\mathbf{T}^{E^-}})]_{\text{virt}} \\ &\in H^0(D^{E^-} \times \{\text{pt}_{\mathbf{T}^{E^-}}\}, \mathbb{Q}) = \mathbb{Q} \end{aligned}$$

where

$$\begin{aligned} \mathcal{M}_{1,1}^{\text{op}}(\beta_j + \alpha; D_i; \text{pt}_{\mathbf{T}^X}) &= (\mathcal{M}_{1,1}^{\text{op}}(\beta_j + \alpha) \times_X D_i) \times_{\mathbf{T}^X} \{\text{pt}_{\mathbf{T}^X}\} \\ &= \mathcal{M}_{1,1}^{\text{op}}(\beta_j + \alpha) \times_{X \times \mathbf{T}^X} (D_i \times \{\text{pt}_{\mathbf{T}^X}\}) \end{aligned}$$

and

$$\begin{aligned} \mathcal{M}_{1,1}^{\text{op}}(b_0 + b_j + \alpha; D_i^{E^-}; \text{pt}_{\mathbf{T}^{E^-}}) &= (\mathcal{M}_{1,1}^{\text{op}}(b_0 + b_j + \alpha) \times_{E^-} D_i^{E^-}) \times_{\mathbf{T}^{E^-}} \{\text{pt}_{\mathbf{T}^{E^-}}\} \\ &= \mathcal{M}_{1,1}^{\text{op}}(b_0 + b_j + \alpha) \times_{E^- \times \mathbf{T}^{E^-}} (D_i^{E^-} \times \{\text{pt}_{\mathbf{T}^{E^-}}\}). \end{aligned}$$

The fiber products appeared above use the evaluation maps $\text{ev}_+^X : \mathcal{M}_{1,1}^{\text{op}}(\beta_j + \alpha) \rightarrow X$, $\text{ev}_0^X : \mathcal{M}_{1,1}^{\text{op}}(\beta_j + \alpha) \rightarrow \mathbf{T}^X$, $\text{ev}_+^{E^-} : \mathcal{M}_{1,1}^{\text{op}}(\beta_j + \alpha) \rightarrow E^-$, $\text{ev}_0^{E^-} : \mathcal{M}_{1,1}^{\text{op}}(\beta_j + \alpha) \rightarrow \mathbf{T}^{E^-}$, and the inclusion maps $D_i \hookrightarrow X$, $\{\text{pt}_{\mathbf{T}^X}\} \hookrightarrow \mathbf{T}^X$, $D_i^E \hookrightarrow E^-$, $\{\text{pt}_{\mathbf{T}^{E^-}}\} \hookrightarrow \mathbf{T}^{E^-}$.

Thus to prove

$$n_{1,1}^X(\beta_j + \alpha; D; [\text{pt}]_{\mathbf{T}^X}) = n_{1,1}^{E^-}(b_0 + b_j + \alpha; D^{E^-}; [\text{pt}]_{\mathbf{T}^{E^-}})$$

it suffices to show that

$$\mathcal{M}_{1,1}^{\text{op}}(\beta_j + \alpha; D; \text{pt}_{\mathbf{T}^X}) \cong \mathcal{M}_{1,1}^{\text{op}}(b_0 + b_j + \alpha; D^{E^-}; \text{pt}_{\mathbf{T}^{E^-}})$$

as Kuranishi spaces. This is the content of the next proposition.

Proposition 5.5. *Fix a point $\text{pt}_{\mathbf{T}^X} \in \mathbf{T}^X$ and a point $\text{pt}_{\mathbf{T}^{E^-}} \in \mathbf{T}^{E^-}$. Then we have*

$$(5.2) \quad \mathcal{M}_{1,1}^{\text{op}}(\beta_j + \alpha; D; \text{pt}_{\mathbf{T}^X}) \cong \mathcal{M}_{1,1}^{\text{op}}(b_0 + b_j + \alpha; D^{E^-}; \text{pt}_{\mathbf{T}^{E^-}})$$

as Kuranishi spaces.

Proof. We divide the proof into three parts.

(A) Virtual dimensions. First of all, both sides have virtual dimension zero:

$$\dim \mathcal{M}_{1;1}^{\text{op}}(\beta_j + \alpha) = \mu(\beta_j) + 2c_1(\alpha) + 2 + 1 + n - 3 = 2 + n.$$

Requiring the interior marked point to pass through D cuts down the dimension by 2; Requiring the boundary marked point to pass through $\text{pt}_{\mathbf{T}^X}$ further cuts down the dimension by n . Thus the virtual dimension of the LHS of (5.2) is zero. For the RHS of (5.2),

$$\dim \mathcal{M}_{1;1}^{\text{op}}(b_0 + b_j + \alpha) = \mu(b_0) + \mu(b_j) + 2c_1(\alpha) + 2 + 1 + (n + 1) - 3 = 5 + n.$$

Requiring the interior marked point to pass through D^{E^-} cuts down the dimension by 4; Requiring the boundary marked point to pass through $\text{pt}_{\mathbf{T}^{E^-}}$ further cuts down the dimension by $n + 1$. Thus the virtual dimension of the RHS of (5.2) is also zero.

(B) Spaces. In what follows the domain interior marked point of a stable disc is always denoted as p^{int} , and the domain boundary marked point is always denoted as p^{bdy} .

Now we construct a bijection between the left hand side and the right hand side of (5.2). In the following we fix a local toric chart $\chi = (\chi_1, \dots, \chi_n)$ of X which covers the open orbit of $D_j \subset X$, and such that $\chi_1(D_j) = 0$. Without loss of generality, we may take $\mathbf{T}^X$ to be the fiber $|\chi_l| = 1$ for all l , and $\text{pt}_{\mathbf{T}^X}$ to be $\chi_l(\text{pt}_{\mathbf{T}^X}) = 1$ for all l . Correspondingly we have the local chart (χ, w) of E^- around the fiber $w = 0 \in \mathbb{P}^1$. Without loss of generality we take $\mathbf{T}^{E^-}$ to be the fiber $|\chi_l| = |w| = 1$, and $\text{pt}_{\mathbf{T}^{E^-}}$ to be $\chi_l(\text{pt}_{\mathbf{T}^{E^-}}) = w(\text{pt}_{\mathbf{T}^{E^-}}) = 1$ for all l .

First consider the easier case $\alpha = 0$. By Lemma 2.3, a stable disc in the LHS is a holomorphic disc u in class β_j . The domain is a closed unit disc $\Delta \in \mathbb{C}$. By using automorphism we may take $p^{\text{int}} = 0$ and $p^{\text{bdy}} = 1$. Under the above chosen local coordinates of X , u has the expression

$$u(z) = (\mathbf{e}^{i\theta_1} \frac{z - \alpha_0}{1 + \bar{\alpha}_0 z}, \mathbf{e}^{i\theta_2}, \dots, \mathbf{e}^{i\theta_n})$$

for some $\alpha_0 \in \Delta$, $\theta_k \in \mathbb{R}$, $k = 1, \dots, n$. u passes through $D = D_i$ only when $i = j$. Thus the left hand side is simply an empty set when $i \neq j$. When $i = j$, $u(0) \in D_j$ forces $\alpha_0 = 0$, and requiring $u(1) = \text{pt}_{\mathbf{T}^X} = (1, \dots, 1)$ fixes $\theta_k \cong 0$ for all $k = 1, \dots, n$. Thus the left hand side is simply an empty set when $i \neq j$, and is a singleton when $i = j$. On the other side, by Lemma 5.4, a stable disc in the RHS is a holomorphic disc ν in class $b_0 + b_j$. Such disc is also classified by Cho-Oh [10]. Again we use the domain automorphism to fix $p^{\text{bdy}} = 1$ and $p^{\text{int}} = 0$. Then the disc is of the form

$$w \circ \nu(z) = \mathbf{e}^{i\theta_0} \frac{z - \alpha_1}{1 + \bar{\alpha}_1 z}, \quad \chi \circ \nu(z) = (\mathbf{e}^{i\theta_1} \frac{z - \alpha_2}{1 + \bar{\alpha}_2 z}, \mathbf{e}^{i\theta_2}, \dots, \mathbf{e}^{i\theta_n}),$$

where $\alpha_1, \alpha_2 \in \Delta$ and $\theta_i \in \mathbb{R}$. ν never hits $D_i^{E^-}$ when $i \neq j$. When $i = j$, $\nu(0) \in D_j^{E^-}$ forces $w = \chi_1 = 0$ when $z = 0$. Then $\alpha_1 = \alpha_2 = 0$. Also $\nu(1) = \text{pt}_{\mathbf{T}^{E^-}}$ means $w = \chi_1 = \dots = \chi_n = 1$ when $z = 1$, which implies $\theta_0 = \dots = \theta_n = 1$. Thus the moduli space in the RHS is empty when $i \neq j$, and is a singleton when $i = j$. This verifies that the LHS matches with the RHS.

Now consider the case $\alpha \neq 0$. Let ϕ be a stable disc bounded by $\mathbf{T}^X$ in the LHS of (5.2). We associate ϕ with a stable disc bounded by $\mathbf{T}^{E^-}$ in the RHS of (5.2) as follows. By Lemma 2.3, ϕ is a holomorphic disc in class β_j attached with a rational curve in class α at an interior

nodal point. Let us identify the domain disc component with the closed unit disc $\Delta \subset \mathbb{C}$, denote the domain of the rational curve by C , and denote $\phi_\Delta := \phi|_\Delta$, $\phi_C := \phi|_C$. The nodal point corresponds to a point $p^{\text{nod}} \in C$ and a point in Δ . By using automorphism of Δ we may assume this point to be 0 and $p^{\text{bdy}} = 1$. Then $\phi_\Delta(0) = \phi_C(p^{\text{nod}})$. Under the chosen local coordinates χ , we have

$$\phi_\Delta(z) = (e^{i\theta_1} \frac{z - \alpha_0}{1 + \bar{\alpha}_0 z}, e^{i\theta_2}, \dots, e^{i\theta_n})$$

for some $\alpha_0 \in \Delta$, $\theta_l \in \mathbb{R}$ for $l = 1, \dots, n$.

Since ϕ_C has Chern number zero, $\phi(C) \subset \bigcup_l D_l$, and in particular $\phi_\Delta(0) = \phi_C(p^{\text{nod}}) \in \bigcup_l D_l$. But ϕ_Δ does not hit any toric divisors except D_j . Thus $\phi_\Delta(0) \in D_j$, and $0 \in \Delta$ is the only point which is mapped to $\bigcup_l D_l$ under ϕ_Δ . This forces $\alpha_0 = 0$ in the above expression of ϕ_Δ . Moreover ϕ_Δ maps $z = 1$ to $\text{pt}_{\mathbf{T}^x} = (1, \dots, 1)$, and this forces $\theta_1 \cong \theta_2 \cong \dots \cong \theta_n \cong 0$. As a result, $\phi_\Delta = (z, 1, \dots, 1)$.

On the other hand $\phi(p^{\text{int}}) \in D_i$. Suppose p^{int} lies on the disc component. Since p^{int} has to be different from the nodal point, $p^{\text{int}} \neq 0$. But then $\phi_\Delta(p^{\text{int}}) \notin \bigcup_l D_l$, and so p^{int} is not mapped to D_i , a contradiction. Thus p^{int} has to be located in the rational curve C .

We associate to ϕ an element ϕ^{E^-} in the RHS which has the same domain and marked points $p^{\text{bdy}}, p^{\text{int}}$ as ϕ (the domain is Δ attached with C at $z = 0$). $\phi^{E^-}|_\Delta$ is defined to be $(\phi_\Delta(z), z)$ written in terms of the above chosen local coordinates (χ, w) of E^- , and $\phi^{E^-}|_C := (\phi_C, 0)$. Notice that $\phi^{E^-}|_\Delta(0) = (\phi_\Delta(0), 0) = (\phi_C(p^{\text{nod}}), 0) = \phi^{E^-}|_C(p^{\text{nod}})$, and so ϕ^{E^-} is well-defined. Moreover since $\phi_C(p^{\text{int}}) \in D_i$, $\phi^{E^-}(p^{\text{int}}) = (\phi_C(p^{\text{int}}), 0) \in D_i^{E^-}$. Also $\phi^{E^-}(p^{\text{bdy}}) = (\phi_\Delta(1), 1) = \text{pt}_{\mathbf{T}^{E^-}}$. This verifies that ϕ^{E^-} is an element in the RHS.

Now we prove that every element in the RHS of (5.2) comes from an element from the LHS of (5.2) in the way described above. By Lemma 5.4, a stable disc ϕ^{E^-} in $b_0 + b_j + \alpha$ must be a holomorphic disc representing $b_0 + b_j$ attached with a rational curve of Chern number zero representing α . As above, the domain disc component is identified with the unit disc $\Delta \subset \mathbb{C}$, and the domain rational curve is denoted by C . The nodal point corresponds to a point $p^{\text{nod}} \in C$ and a point in Δ . By using automorphism of Δ we may assume this point to be 0 and $p^{\text{bdy}} = 1$. Then $\phi_\Delta^{E^-}(0) = \phi_C^{E^-}(p^{\text{nod}})$. Using Cho-Oh's classification of holomorphic discs [10], in the chosen local coordinates (w, χ) , $\phi_\Delta^{E^-}$ is of the form

$$w \circ \phi_\Delta^{E^-}(z) = e^{i\theta_0} \frac{z - \alpha_0}{1 + \bar{\alpha}_0 z}, \quad \chi \circ \phi_\Delta^{E^-}(z) = (e^{i\theta_1} \frac{z - \alpha_1}{1 - \bar{\alpha}_1 z}, e^{i\theta_2}, \dots, e^{i\theta_n}).$$

Suppose p^{int} lies in the disc component. Then $\phi_\Delta^{E^-}(p^{\text{int}}) \in D_i^{E^-}$. This happens only when $i = j$, $\alpha_0 = \alpha_1 = 0$. In such case $\phi_\Delta^{E^-}$ hits the union of toric divisors of E^- only at one point $z = p^{\text{int}}$. Now $\phi_C^{E^-}$ represents the fiber class α with $c_1(\alpha) = 0$, and so by Lemma 5.2 $\phi^{E^-}(C) \subset \bigcup_{l=1}^m D_l$. In particular $\phi_\Delta^{E^-}(0) = \phi_C^{E^-}(p^{\text{nod}}) \in \bigcup_{l=1}^m D_l$. This forces p^{int} to coincide with the nodal point, a contradiction. Thus p^{int} must lie in the rational curve C .

The image of $\phi_C^{E^-}$ lies in a fiber of $E^- \rightarrow \mathbb{P}^1$. But since $\phi_C^{E^-}(p^{\text{int}}) \in D_i^{E^-}$ which lies in $\mathcal{D}_0$ (the fiber at zero), this forces $\phi_C^{E^-}$ to lie in $\mathcal{D}_0$. Then $\phi_C^{E^-}$ is of the form $(0, \phi_C)$ in the local coordinates (w, χ) . Together with $\phi^{E^-}(C) \subset \bigcup_{l=1}^m D_l$, this means $\phi^{E^-}(C) \subset \bigcup_{l=1}^m D_l^{E^-}$. Then $\phi_\Delta^{E^-}(0) = \phi_C^{E^-}(p^{\text{nod}}) \in \bigcup_{l=1}^n D_l^{E^-}$, which happens only when $\alpha_0 = \alpha_1 = 0$. Moreover $\phi_\Delta^{E^-}(1) = (1, \dots, 1)$, and so $\theta_0 = \dots = \theta_n = 1$. Thus $\phi_\Delta^{E^-} = (z, \phi_\Delta(z))$ in the local coordinates

(w, χ) , where $\phi_\Delta(z) = (z, 1, \dots, 1)$. Thus ϕ^{E^-} comes from the stable disc ϕ in X , which is a union of ϕ_Δ and ϕ_C .

(C) Kuranishi Structures. Now we compare the Kuranishi structures on the both sides of (5.2). Let us have a brief reasoning on why they should have the same Kuranishi structures. On both sides the disc components are regular, and so the obstructions merely come from the rational curve components in class α . For the curve component of ϕ^{E^-} , since it is free to move from fiber to fiber of $E^- \rightarrow \mathbb{P}^1$, the obstruction comes from the directions along X , and this is identical with the corresponding curve component of ϕ . Now consider the deformations. Due to the boundary point condition, the disc components on both sides cannot be deformed. For the curve component of ϕ^{E^-} , the interior point condition that it has to pass through D^{E^-} kills the deformations in the direction transverse to fibers. Thus ϕ^{E^-} has the same deformations as ϕ . Thus the corresponding stable discs on both sides have the same deformations and obstructions, and hence the moduli have the same Kuranishi structures. In what follows, we write down and equate the deformations and obstructions on both sides.

A Kuranishi structure on $\mathcal{M}_{1,1}^{\text{op}}(\beta_j + \alpha; D; \text{pt}_{\mathbf{T}^X})$ assigns a Kuranishi chart

$$(V_{\text{op}}, \mathcal{E}_{\text{op}}^-, \Gamma_{\text{op}}, \psi_{\text{op}}, s_{\text{op}})$$

around each $\phi \in \mathcal{M}_{1,1}^{\text{op}}(\beta_j + \alpha; D; \text{pt}_{\mathbf{T}^X})$ which is constructed as follows. Let

$$D_\phi \bar{\partial} : W^{1,p}(\text{Dom}(\phi), \phi^*(TX), \mathbf{T}) \rightarrow W^{0,p}(\text{Dom}(\phi), u^*(TX) \otimes \Lambda^{0,1})$$

be the linearized Cauchy-Riemann operator at ϕ . (Here $\text{Dom}(\phi)$ is the domain of ϕ .)

- (1) Γ_{op} is the automorphism group of ϕ , that is, the group of all elements

$$g \in \text{Aut}(\text{Dom}(\phi), p^{\text{int}}, p^{\text{bdy}})$$

such that $\phi \circ g = \phi$. By stability of ϕ , Γ_{op} is a finite group. (Note that by definition, $g(p^{\text{int}}) = p^{\text{int}}, g(p^{\text{bdy}}) = p^{\text{bdy}}$.)

- (2) The so-called obstruction space $\mathcal{E}_{\text{op}}^-$ is the cokernel of the linearized Cauchy-Riemann operator $D_\phi \bar{\partial}$, which is finite dimensional since $D_\phi \bar{\partial}$ is Fredholm. For the purpose of the next step of construction, it is identified (in a non-canonical way) with a subspace of $W^{0,p}(\text{Dom}(\phi), \phi^*(TX) \otimes \Lambda^{0,1})$ as follows. Denote by Δ and $S_1, \dots, S_l$ the disc and sphere components of $\text{Dom}(\phi)$ respectively. Take non-empty open subsets $W_0 \subset \Delta$ and $W_i \subset S_i$ for $i = 1, \dots, l$. Then by unique continuation theorem there exists finite dimensional subspaces $\mathcal{E}_i^- \subset C_0^\infty(W_i, \phi^*(TX) \otimes \Lambda^{0,1})$ such that

$$\text{Im}(D_\phi \bar{\partial}) \oplus \mathcal{E}_{\text{op}}^- = W^{0,p}(\text{Dom}(\phi), \phi^*(TX) \otimes \Lambda^{0,1})$$

and $\mathcal{E}_{\text{op}}^-$ is invariant under Γ_{op} , where

$$\mathcal{E}_{\text{op}}^- := \mathcal{E}_0^- \oplus \dots \oplus \mathcal{E}_l^-.$$

- (3) $\tilde{V}_{\text{op}}$ is taken to be (a neighborhood of 0 of) the space of first order deformations Φ of ϕ which satisfies the linearized Cauchy-Riemann equation modulo elements in $\mathcal{E}_{\text{op}}^-$, that is,

$$(D_\phi \bar{\partial}) \cdot \Phi \equiv 0 \quad \text{mod } \mathcal{E}_{\text{op}}^-.$$

Such deformations may come from deformations of the map or deformations of complex structures of the domain. More precisely,

$$\tilde{V}_{\text{op}} = V_{\text{op}}^{\text{map}} \times V_{\text{op}}^{\text{dom}}$$

where $V_{\text{op}}^{\text{map}}$ is defined in the following way. Let $V'_{\text{op, map}}$ be the kernel of the linear map

$$[D_{\phi}\bar{\partial}] : W^{1,p}(\text{Dom}(\phi), \phi^*(TX), \mathbf{T}) \rightarrow W^{0,p}(\text{Dom}(\phi), \phi^*(TX) \otimes \Lambda^{0,1})/\mathcal{E}_{\text{op}}^-.$$

Notice that $\text{Aut}(\text{Dom}(\phi), p^{\text{int}}, p^{\text{bdy}})$ may not be finite since the domain of ϕ may not be stable, and it acts on $V'_{\text{op, map}}$. Thus its Lie algebra $\mathfrak{g}$ is contained in $V'_{\text{op, map}}$, and we take $V_{\text{op}}^{\text{map}} \subset V'_{\text{op, map}}$ such that $V'_{\text{op, map}} = V_{\text{op}}^{\text{map}} \oplus \mathfrak{g}$.

$V_{\text{op}}^{\text{dom}}$ is a neighborhood of zero in the space of deformations of the domain rational curve C . Such deformations consists of two types: one is deformations of each stable component (in this genus zero case, it means movements of special points in each component), and another one is smoothing of nodes between components. That is,

$$V_{\text{op}}^{\text{dom}} = V_{\text{op}}^{\text{cpnt}} \times V_{\text{op}}^{\text{smth}}$$

where $V_{\text{op}}^{\text{cpnt}}$ is a neighborhood of zero in the space of deformations of components of C , and $V_{\text{op}}^{\text{smth}}$ is a neighborhood of zero in the space of smoothing of the nodes (each node contributes to a one-dimensional family of smoothings). Each deformation in $V_{\text{op}}^{\text{dom}}$ gives $\Delta \cup \tilde{C}$, where Δ is a disc with one boundary marked point, and $\tilde{C}$ is a rational curve with one interior marked point, such that Δ and $\tilde{C}$ intersect at a nodal point. $\Delta \cup \tilde{C}$ serves as the domain of the deformed map Φ .

- (4) $\tilde{s}_{\text{op}} : \tilde{V}_{\text{op}} \rightarrow \mathcal{E}_{\text{op}}^-$ is a transversal Γ_{op} -equivariant perturbed zero-section of the trivial bundle $\mathcal{E}_{\text{op}}^- \times \tilde{V}_{\text{op}}$ over $\tilde{V}_{\text{op}}$. By [14], this can be chosen to be $\mathbf{T}$ -equivariant.
- (5) There exists a continuous family of smooth maps $\rho_{\Phi}^{\text{op}} : (\mathcal{D}, \partial\mathcal{D}) \rightarrow (X, \mathbf{T})$ over $\tilde{V}_{\text{op}} \ni \Phi$ such that it solves the inhomogeneous Cauchy-Riemann equation: $\bar{\partial}\rho_{\Phi}^{\text{op}} = \tilde{s}_{\text{op}}(\Phi)$. Set

$$V_{\text{op}} := \{\Phi \in \tilde{V}_{\text{op}} : \text{ev}_0(\rho_{\Phi}^{\text{op}}) = \text{pt}_{\mathbf{T}^X}; \text{ev}_+(\rho_{\Phi}^{\text{op}}) \in D\}$$

where ev_0 is the evaluation map at p^{bdy} . Then set $s_{\text{op}} := \tilde{s}_{\text{op}}|_{V_{\text{op}}}$.

- (6) ψ_{op} is a map from $s_{\text{op}}^{-1}(0)/\Gamma_{\text{op}}$ onto a neighborhood of $[\phi] \in \mathcal{M}_{1,1}^{\text{op}}(\beta_j + \alpha; D; \text{pt}_{\mathbf{T}^X})$.

Now comes the key: in Item (2) of the above construction, since the disc component of ϕ is unobstructed (that is, the linearized Cauchy-Riemann operator localized to the disc component is surjective), $\mathcal{E}_0^- = 0$ so that $\mathcal{E}_{\text{op}}^-$ is of the form

$$\mathcal{E}_{\text{op}}^- = 0 \oplus \mathcal{E}_1^- \oplus \dots \oplus \mathcal{E}_l^-.$$

The analogous statement is also true for the corresponding stable disc $\phi^{E^-} \in \mathcal{M}_{1,1}^{\text{op}}(b_0 + b_j + \alpha; D^{E^-}; \text{pt}_{\mathbf{T}^{E^-}})$. With this observation, we argue in the following that $(V_{\text{op}}, \mathcal{E}_{\text{op}}^-, \Gamma_{\text{op}}, \psi_{\text{op}}, s_{\text{op}})$ can be identified as a Kuranishi chart $(V_{\text{op}}^E, \mathcal{E}_{\text{op}}^{-,E}, \Gamma_{\text{op}}^E, \psi_{\text{op}}^E, s_{\text{op}}^E)$ around the corresponding stable disc ϕ^{E^-} bounded by $\mathbf{T}^{E^-} \subset E^-$.

- (1) ϕ and ϕ^{E^-} have the same automorphism group, that is, $\Gamma_{\text{op}} = \Gamma_{\text{op}}^E$. This is because the disc component have only one boundary marked point and one interior nodal point and thus has no automorphism, and any automorphism on the rational-curve part of ϕ will give an automorphism on the rational-curve part of ϕ^{E^-} , and vice versa.

- (2) The disc component of ϕ^{E^-} is unobstructed. For the rational curve component C which is mapped into $\mathcal{D}_0 \cong X$, notice that there is a splitting $TE|_{\mathcal{D}_0} = T(\mathcal{D}_0) \oplus N\mathcal{D}_0$ and so $W^{0,p}(C, (\phi^{E^-}|_C)^*(TE) \otimes \Lambda^{0,1})$ is equal to

$$W^{0,p}(C, (\phi^{E^-}|_C)^*(T\mathcal{D}_0) \otimes \Lambda^{0,1}) \oplus W^{0,p}(C, (\phi^{E^-}|_C)^*(N\mathcal{D}_0) \otimes \Lambda^{0,1})$$

where the first summand is equal to $W^{0,p}(C, (\phi)^*TX \otimes \Lambda^{0,1})$.

Since the curve component is free to move in the direction of the normal bundle $N\mathcal{D}_0$, we have

$$\text{Im}(D_{\phi^{E^-}|_C} \bar{\partial}) \supset W^{0,p}(C, (\phi^{E^-}|_C)^*(N\mathcal{D}_0) \otimes \Lambda^{0,1}).$$

Hence

$$\text{Im}(D_{\phi^{E^-}} \bar{\partial}) \oplus (0 \oplus \mathcal{E}_1^- \oplus \dots \oplus \mathcal{E}_l^-) = W^{0,p}(\text{Dom}(\phi^{E^-}), (\phi^{E^-})^*(TX) \otimes \Lambda^{0,1}).$$

Thus we may take $\mathcal{E}_{\text{op}}^{-,E} = 0 \oplus \mathcal{E}_1^- \oplus \dots \oplus \mathcal{E}_l^-$.

- (3) $\tilde{V}_{\text{op}}^E$, $\tilde{s}_{\text{op}}^E$, $\rho_{\Phi^{E^-}}^{E^-, \text{op}}$ are defined in the same way as above. The subspace $\tilde{V}'$ of those deformations $\Phi^{E^-} \in \tilde{V}_{\text{op}}^E$ such that the image of the curve component under $\rho_{\Phi^{E^-}}^{E^-, \text{op}}$ lies in $\mathcal{D}_0$ is isomorphic to $\tilde{V}_{\text{op}}$, and restrictions of $\tilde{s}_{\text{op}}^E$ and $\rho_{\Phi^{E^-}}^{E^-, \text{op}}$ to $\tilde{V}'$ gives choices of $\tilde{s}_{\text{op}}$ and ρ^{op} respectively. Moreover,

$$V_{\text{op}}^E := \{\Phi^{E^-} \in \tilde{V}_{\text{op}}^E : \text{ev}_0(\rho_{\Phi^{E^-}}^{E^-, \text{op}}) = \text{pt}_{\mathbf{T}^{E^-}}; \text{ev}_+(\rho_{\Phi^{E^-}}^{E^-, \text{op}}) \in D^{E^-}\}$$

lies in $\tilde{V}_{\text{op}}$. Thus $V_{\text{op}}^E = V_{\text{op}}$, and $s_{\text{op}}^E := \tilde{s}_{\text{op}}^E|_{V_{\text{op}}^E} = s_{\text{op}}$. Then ψ_{op} can be identified as a map ψ_{op}^E which maps $(s_{\text{op}}^E)^{-1}(0)/\Gamma_{\text{op}}^E$ onto a neighborhood of $[\phi^{E^-}] \in \mathcal{M}_{1,1}^{\text{op}}(b_0 + b_j + \alpha; D^{E^-}; \text{pt}_{\mathbf{T}^{E^-}})$.

In conclusion, a Kuranishi neighborhood of ϕ can be identified with a Kuranishi neighborhood of ϕ^{E^-} . Thus the Kuranishi structures on $\mathcal{M}_{1,1}^{\text{op}}(\beta; D; \text{pt}_{\mathbf{T}^X})$ and that on $\mathcal{M}_{1,1}^{\text{op}}(b_0 + b_j + \alpha; D^{E^-}; \text{pt}_{\mathbf{T}^{E^-}})$ are identical. This completes the proof of the proposition. $\square$

5.2. Second step. Now we come to the second main step, which is the following theorem:

Theorem 5.6. *Assume the notations as in Theorem 5.1. Then*

$$n_{1,1}^{E^-}(b_0 + b_j + \alpha; D^{E^-}; [\text{pt}]_{\mathbf{T}^{E^-}}) = \langle D^{E^-}, [\text{pt}]_{E^-} \rangle_{0,2,\sigma^-+\alpha}^{E^-}.$$

By Equation (5.1),

$$n_{1,1}^{E^-}(b_0 + b_j + \alpha; D^{E^-}; [\text{pt}]_{\mathbf{T}^{E^-}}) = [\mathcal{M}_{1,1}^{\text{op}}(b_0 + b_j + \alpha; D^{E^-}; \text{pt}_{\mathbf{T}^{E^-}})]_{\text{virt}} \in H^0(D^{E^-} \times \{\text{pt}_{\mathbf{T}^{E^-}}\}, \mathbb{Q}) = \mathbb{Q}$$

and

$$\langle D^{E^-}, [\text{pt}]_{E^-} \rangle_{0,2,\sigma^-+\alpha}^{E^-} = [\mathcal{M}_{0,2,\sigma^-+\alpha}^{\text{cl}}(D^{E^-}; \text{pt})]_{\text{virt}} \in H^0(\text{pt}, \mathbb{Q}) = \mathbb{Q}$$

where $\mathcal{M}_{0,2,b_0+b_j+\alpha}^{\text{cl}}(D^{E^-}; \text{pt})$ is the fiber product $\mathcal{M}_{0,2,b_0+b_j+\alpha}^{\text{cl}} \times_{E^- \times E^-} (D^{E^-} \times \text{pt})$ for any chosen point $\text{pt} \in E^-$. In order to prove the equality between open and closed invariants in Theorem 5.6, it suffices to show an isomorphism between the Kuranishi structures stated as follows:

Proposition 5.7. *Fix a point $\text{pt} \in \mathbf{T}^{E^-} \subset E^-$. Then*

$$(5.3) \quad \mathcal{M}_{1;1}^{\text{op}}(b_0 + b_j + \alpha) \times_{E^- \times \mathbf{T}^{E^-}} (D_i^{E^-} \times \{\text{pt}\}) \cong \mathcal{M}_{0,2,\sigma^-+\alpha}^{\text{cl}} \times_{E^- \times E^-} (D_i^{E^-} \times \{\text{pt}\}).$$

as Kuranishi spaces.

The proof is very similar to that of Proposition 5.5: We first prove that the two sides are equal as sets, and then compare the Kuranishi charts and show that they can be chosen to be the same. First, let us consider the case $\alpha = 0$ and $i \neq j$, in which the invariants become zero:

Proof of Proposition 5.7 when $\alpha = 0$ and $i \neq j$. We have seen in the proof of Proposition 5.5 that the LHS of (5.3) is an empty set when $\alpha = 0$ and $i \neq j$.

For the right hand side, the only possibility for a rational curve in σ^- to pass through $D_i^{E^-}$ and pt is that it degenerates to the union of a curve representing σ_0^- and a fiber curve. Such a curve is unique since pt fixes the location of the fiber curve, and by connectedness the location of the curve in σ_0^- is fixed. Thus the moduli is a singleton. This moduli has a non-trivial obstruction bundle, which is the negative normal bundle of the curve in σ_0^- . Thus as a Kuranishi space the moduli is equivalent to an empty set. In this case $n_{1,1}^{E^-}(b_0 + b_j; D_j^{E^-}; [\text{pt}]_{\mathbf{T}^{E^-}}) = \langle D_j^{E^-}, [\text{pt}]_{E^-} \rangle_{0,2,\sigma^-}^{E^-} = 0$. $\square$

Proof of Proposition 5.7 when $\alpha \neq 0$ or $i = j$. First we construct a bijection between the two sides of (5.3).

For a stable disc in $\mathcal{M}_{1;1}^{\text{op}}(b_0 + b_j + \alpha) \times_{E^- \times \mathbf{T}^{E^-}} (D_i^{E^-} \times \{\text{pt}\})$, we denote the domain interior marked point by p^{int} and the domain boundary marked point by p^{bdy} . For a rational curve in $\mathcal{M}_{0,2,\sigma^-+\alpha}^{\text{cl}} \times_{E^- \times E^-} (D_i^{E^-} \times \{\text{pt}\})$, we denote by p_0 the marked point mapped to $D_i^{E^-}$, and p_1 the marked point mapped to pt .

By relabeling $\{D_l\}_{l=1}^m$ if necessary, we assume $j = 1$. Let us fix a local toric chart $\chi = (\chi_1, \dots, \chi_n)$ of X which covers the open orbit of $D_1 \subset X$, and such that $\chi_l|_{D_l} = 0$ (by relabeling $\{D_l\}_{l=1}^m$ if necessary). Correspondingly we have the local chart (χ, w) of E^- around the fiber $w = 0 \in \mathbb{P}^1$. Without loss of generality we take $\mathbf{T}^{E^-}$ to be the fiber $|\chi_l| = |w| = 1$ for all l , and $\text{pt} \in \mathbf{T}^{E^-}$ to be $\chi_l(\text{pt}) = w(\text{pt}) = 1$ for all l .

Consider the case $\alpha = 0$ and $i = j$. We have seen in the proof of Proposition 5.5 that the LHS of (5.3) is a singleton when $i = j$. When $i = j$, it is the disc $w = z, \chi = (z, 1, \dots, 1)$ on $\Delta \ni z$.

On the RHS of (5.3), the elements are holomorphic spheres $\rho : \mathbb{P}^1 \rightarrow E^-$ representing σ^- passing through pt and $D_i^{E^-}$. Since $\sigma^- \cdot \mathcal{D}_\infty = 1$, there is a unique point $p_\infty \in \mathbb{P}^1$ with $\rho(p_\infty) \in \mathcal{D}_\infty$. By composing with an automorphism of $\mathbb{P}^1$, we may assume $p_0 = 0, p_1 = 1$ and $p_\infty = \infty$. Consider $\chi_l \circ \rho, w \circ \rho : \mathbb{P}^1 \rightarrow \mathbb{P}^1$ for $l = 1, \dots, n$. Since $\sigma^- \cdot \mathcal{D}_l = 0$ except when $l = 0, j, \infty$, $\chi_l \circ \rho$ are constants for $l = 2, \dots, n$. Thus $\chi_l \circ \rho = \chi_l \circ \rho(p_1) = 1$ for $l = 2, \dots, n$. Moreover $\chi_1 \circ \rho(z)$ and $w \circ \rho(z)$ have only one zero at 0 and one pole at ∞ , and so they are equal to cz for some $c \in \mathbb{C}$. But $\rho(p_1) = \text{pt}$ implies $\chi_1 \circ \rho(p_1) = 1$, and this forces $c = 1$. Thus ρ is $(w, \chi) = (z, (z, 1, \dots, 1))$. This proves that the moduli in the RHS consists of a singleton. Thus there is a bijection between the LHS and RHS of (5.3).

Now consider the case $\alpha \neq 0$. Let $\phi_{\text{op}}^{E^-}$ be a stable disc in the LHS. From the proof of Proposition 5.5, $\phi_{\text{op}}^{E^-}$ is a holomorphic disc representing $b_0 + b_j$ attached with a rational curve C representing α at exactly one nodal point, where the interior marked point p^{int} is located in C , and the map $\phi_{\text{op}}^{E^-}|_{\Delta}$ on the disc component $\Delta \ni z$ is given by $(w, \chi) = (z, (z, 1, \dots, 1))$. Such a map from Δ to E^- analytically extends to a map $\phi_{\text{cl}, \mathbb{P}^1} : \mathbb{P}^1 \rightarrow E^-$, where $\infty \in \mathbb{P}^1$ is mapped to $w = \infty$ and $\chi_l = 1$ for $l = 2, \dots, n$, which is the point $\text{pt} \in \mathcal{D}_{\infty}$. Then $\phi_{\text{cl}, \mathbb{P}^1}$ attached with the same rational curve C , with marked points $p_0 = p^{\text{int}}$ in the rational curve and $p_1 = \infty \in \text{Dom}(\phi_{\text{cl}, \mathbb{P}^1})$, is an element in the moduli on the RHS. This gives a map from the LHS to the RHS of (5.3).

Now we show that this map is invertible. An element in the RHS is a union of a holomorphic sphere $\rho_{\mathbb{P}^1} : \mathbb{P}^1 \rightarrow E^-$ representing σ_0^- and a rational curve $\rho_C : C \rightarrow E^-$ representing α . The requirement that it passes through D^{E^-} forces the image of ρ_C to lie in $\bigcup_{l=1}^m (\mathcal{D}_0 \cap \mathcal{D}_l)$. Then by connectedness of the rational curve, $\rho_{\mathbb{P}^1}$ has to pass through $\bigcup_{l=1}^m (\mathcal{D}_0 \cap \mathcal{D}_l)$ and pt . By the above argument, $(w, \chi) \circ \rho_{\mathbb{P}^1}(z) = (z, (z, 1, \dots, 1))$ where $\rho_{\mathbb{P}^1}(0)$ is the nodal point. Then by restricting $\rho_{\mathbb{P}^1}$ to $\Delta \subset \mathbb{P}^1$, we obtain a stable disc in the LHS. This gives the inverse of the above map.

The comparison of Kuranishi structures is very much similar to the proof of Proposition 5.5 and thus omitted. $\square$

6. COMPUTING CLOSED INVARIANTS BY SEIDEL REPRESENTATIONS

In this section we prove Theorems 1.1, 1.2, 1.3, 1.5 as promised in the introduction.

6.1. Calculations. We have equated the open Gromov-Witten invariants appearing in the disc potential of X with certain two-point closed Gromov-Witten invariants in the Seidel spaces E_j^- associated to X . Naïvely one would like to compute the closed invariants by using the toric mirror theorem [18, 27] (see Theorem 3.4). Indeed if one pretends that the infinity section class $\sigma_{\infty}^- \in H_2^{\text{eff}}(E_j^-)$ is not there, it is not difficult to deduce Theorem 6.5 from the toric mirror theorem. However, in general, since the infinity section class $\sigma_{\infty}^- \in H_2^{\text{eff}}(E_j^-)$ may have $c_1(\sigma_{\infty}^-) < 0$, E_j^- is *not* semi-Fano. This is because the $\mathbb{C}^*$ -action induced by $-v_j$ can have a fixed locus whose normal bundle has total weights less than -2 . In such circumstances, the mirror theorem involves Birkhoff factorization and generalized mirror transform, which in practice would be rather difficult (if not impossible) to use to compute the closed invariants.

On the other hand, the Seidel space E_j associated to v_j is always semi-Fano [20] because every fixed locus has total weights not less than -2 (the fixed locus D_j has total weight -1 , and the other one has positive weights). Thus the mirror theorem for E_j does not involve Birkhoff factorization and is much easier to handle. In particular, the normalized Seidel element S_j° corresponding to E_j was computed by González-Iritani [20] and can be explicitly expressed in terms of the Batyrev elements:

Proposition 6.1 ([20]).

$$B_j(\check{q}(q)) = \exp(g_j(\check{q}(q)))S_j^{\circ}(q).$$

By [30], the two-point closed Gromov-Witten invariants in question are encoded in the Seidel elements S_j^- (see Proposition 6.2). Together with the equality $S_j * S_j^- = q^{\sigma_j + \sigma_j^-}$

(Corollary 4.5) coming from the Seidel representations, this gives a computation of the closed Gromov-Witten invariants that we need.

Proposition 6.2 ([30], see also Definition 2.5 of [31]).

$$S_j^- * D_i = \sum_k \sum_{d \in H_2^{\text{sec}}(E_j^-)} q^d \langle \iota_* D_i, \iota_* \Phi_k \rangle_{0,2,d}^{E_j^-} \Phi^k \in H^*(X)$$

where $\iota_* : X \hookrightarrow E_j^-$ is the inclusion as a fiber, $\{\Phi^k\}$ is an additive basis of $H^*(X)$ and $\{\Phi_k\} \subset H^*(X)$ is its Poincaré dual basis.

Proposition 6.3. As elements in $QH^*(X)$, we have

$$D_i = B_i + \sum_l (\hat{D}_i \cdot g_l(\check{q}(q))) B_l.$$

Proof. Recall that

$$\tilde{W}_q^{\text{HV}} = \sum_{l=1}^m \exp(g_l(\check{q}(q))) Z_l.$$

Thus under the isomorphism $QH^*(X) \cong \text{Jac}(W^{\text{HV}})$ in Theorem 3.6, D_i is mapped to

$$\begin{aligned} & \sum_{l=1}^m (\delta_{il} + \hat{D}_i \cdot g_l(\check{q}(q))) \exp(g_l(\check{q}(q))) Z_l \\ &= \exp(g_i(\check{q}(q))) Z_i + \sum_{l=1}^m (\hat{D}_i \cdot g_l(\check{q}(q))) \exp(g_l(\check{q}(q))) Z_l. \end{aligned}$$

By Proposition 3.15, B_l is mapped to $\exp(g_l(\check{q}(q))) Z_l$ under the isomorphism. Hence the element

$$B_i + \sum_{l=1}^m (\hat{D}_i \cdot g_l(\check{q}(q))) B_l \in QH^*(X)$$

has the same image as D_i under the isomorphism, and thus we can conclude that

$$D_i = B_i + \sum_l (\hat{D}_i \cdot g_l(\check{q}(q))) B_l.$$

□

Then for all $i, j = 1, \dots, m$, we have

$$\begin{aligned} q^{\sigma_j + \sigma_j^-} D_i &= (S_j * S_j^-) * D_i \\ &= S_j * (S_j^- * D_i) \\ &= q^{\sigma_j} \exp(-g_j(\check{q}(q))) B_j * \sum_k \sum_{d \in H_2^{\text{sec}}(E_j^-)} q^d \langle \iota_* D_i, \iota_* \Phi_k \rangle_{0,2,d}^{E_j^-} \Phi^k. \end{aligned} \tag{6.1}$$

By Proposition 6.3, the left hand side of the above equation is equal to

$$q^{\sigma_j + \sigma_j^-} B_i + q^{\sigma_j + \sigma_j^-} \sum_l (\hat{D}_i \cdot g_l(\check{q}(q))) B_l.$$

Proposition 6.4. $\{B_i\} \cup \{B_l : g_l \neq 0\}$ is a linearly independent set in $H^*(X, \mathbb{Q})$.

Proof. It can be seen from the fan polytope of X [20]. Since X is semi-Fano, the generators v_l of rays lie on the boundary of the fan polytope, and those v_l with $g_l \neq 0$ are not the vertices of the fan polytope. Since number of vertices is at least $n+1$, the number of v_l 's with $g_l \neq 0$ is no more than $m-n-1$. Moreover the only relations among the B_l 's (regarded as elements in a vector space) are the linear relations, and all of them involve elements outside $\{B_l : g_l \neq 0\}$. Thus $\{B_l : g_l \neq 0\}$ is linearly independent. Every linear relation involves more than two vertices, and hence there is no linear relation involving only B_i and B_l 's with $g_l \neq 0$. $\square$

Now we are ready to prove the following main theorem (which is the same as Theorem 1.1):

Theorem 6.5. For all $j = 1, \dots, m$,

$$\exp(g_j(\check{q}(q))) = \sum_{\alpha \in H_2^{c_1=0}(X)} q^\alpha n_1(\beta_j + \alpha) = 1 + \delta_j(q).$$

Proof. Both sides are equal to 1 when $g_j = 0$. So we suppose that $g_j \neq 0$.

Since $\{B_i\} \cup \{B_l : g_l \neq 0\}$ is a linearly independent set (B_i may coincide with B_l for some l though), we may extend it to a basis $\{\Psi^k\}$ of $H^*(X)$. The action of $B_j = \exp g_j(\check{q}(q)) q^{-\sigma_j} S_j$ by quantum multiplication on $QH^*(X)$ is invertible, and so $\{\Phi^k = B_j^{-1} \Psi^k\}$ is also a basis of $H^*(X)$. The elements corresponding to $\{B_l : g_l \neq 0 \text{ or } l = i\}$ are 1 and $B_j^{-1} B_l$ for $l \neq j$. Taking the dual basis of $\{\Phi^k\}$, the dual partner of $\Phi^0 = 1$ is of the form $\Phi_0 = [\text{pt}] + a$, where $a \in H^{\leq 2n-2}(X)$.

Then

$$\begin{aligned} & B_j * \sum_k \sum_{d \in H_2^{\text{sec}}(E_j^-)} q^d \langle \iota_* D_i, \iota_* \Phi_k \rangle_{0,2,d}^{E_j^-} \Phi^k \\ &= \sum_{d \in H_2^{\text{sec}}(E_j^-)} q^d \langle \iota_* D_i, \iota_* ([\text{pt}] + a) \rangle_{0,2,d}^{E_j^-} B_j + \sum_{k \neq 0} \sum_{d \in H_2^{\text{sec}}(E_j^-)} q^d \langle \iota_* D_i, \iota_* \Phi_k \rangle_{0,2,d}^{E_j^-} \Psi^k. \end{aligned}$$

Comparing the coefficients of B_j on both sides of (6.1), we have

$$q^{\sigma_j^-} \exp(g_j(\check{q}(q))) (\delta_{ij} + \hat{D}_i \cdot g_j(\check{q}(q))) = \sum_{d \in H_2^{\text{sec}}(E_j^-)} q^d \langle \iota_* D_i, \iota_* ([\text{pt}] + a) \rangle_{0,2,d}^{E_j^-}.$$

Notice that the Novikov variable appearing in each term of the left hand side is $q^{\sigma_j^- + \alpha}$ for some $\alpha \in H_2^{c_1=0}(X)$. Thus on the right hand side,

$$\langle \iota_* D_i, \iota_* [\text{pt}] \rangle_{0,2,d}^{E_j^-} + \langle \iota_* D_i, \iota_* a \rangle_{0,2,d}^{E_j^-} \neq 0$$

only when $d = \sigma_j^- + \alpha$ for some $\alpha \in H_2^{c_1=0}(X)$. Since $\deg a \leq 2n-2$, $\langle \iota_* D_i, \iota_* a \rangle_{0,2,\sigma_j^- + \alpha}^{E_j^-} = 0$ by dimension reasons.

In conclusion, we have

$$\begin{aligned} q^{\sigma_j^-} \exp(g_j(\check{q}(q)))(\delta_{ij} + \hat{D}_i \cdot g_j(\check{q}(q))) &= \sum_{\alpha \in H_2^{c_1=0}(X)} q^{\sigma_j^- + \alpha} \langle \iota_* D_i, \iota_* [\text{pt}] \rangle_{0,2,\sigma_j^- + \alpha}^{E_j^-} \\ &= q^{\sigma_j^-} \sum_{\alpha \in H_2^{c_1=0}(X)} q^\alpha n_{1,1}(\beta_j + \alpha; D_i; [\text{pt}]_X) \end{aligned}$$

by applying Theorem 5.1.

This implies that

$$\begin{aligned} \hat{D}_i(q^{\beta_j} \exp(g_j(\check{q}(q)))) &= \sum_{\alpha \in H_2^{c_1=0}(X)} q^{\beta_j + \alpha} n_{1,1}(\beta_j + \alpha; D_i; [\text{pt}]_X) \\ &= \hat{D}_i \sum_{\alpha \in H_2^{c_1=0}(X)} q^{\beta_j + \alpha} n_1(\beta_j + \alpha) \end{aligned}$$

for all i , where the last equality follows from the divisor equation. $\square$

6.2. Corollaries. We now describes some consequences of Theorem 6.5.

Corollary 6.6 (=Theorem 1.3). *The coefficients of the disc potential W^{LF} of a compact semi-Fano toric manifold X are convergent power series in the Kähler parameters q .*

Proof. This follows from Theorem 6.5 and the easy fact that the series g_j and the inverse mirror map $\check{q}(q)$ are convergent. $\square$

Corollary 6.7. *The inverse mirror map $\check{q}(q)$ of a compact semi-Fano toric manifold X is written in terms of the generating functions δ_l of open Gromov-Witten invariants as*

$$\check{q}_k(q) = q_k \prod_{l=1}^m (1 + \delta_l(q))^{(D_l, \Psi_k)} = q_k (1 + \delta_{n+k}(q)) \prod_{p=1}^n (1 + \delta_p(q))^{-(v_{n+k}, \nu_p)}.$$

Proof. By (3.2), we have

$$\check{q}_k(q) = q_k \exp(g^{\Psi_k}(\check{q}(q))) = q_k \prod_{l=1}^m (\exp g_l(\check{q}(q)))^{(D_l, \Psi_k)}$$

and thus the equality follows from Theorem 6.5. Moreover, $\Psi_k = \beta_{n+k} - \sum_{p=1}^n (v_{n+k}, \nu_p) \beta_p$, and so $(D_p, \Psi_k) = -(v_{n+k}, \nu_p)$ for $p = 1, \dots, n$ and $(D_{n+r}, \Psi_k) = \delta_{rk}$ for $r = 1, \dots, m - n$. $\square$

The following Corollary is the same as Theorem 1.2.

Corollary 6.8. *For every semi-Fano toric manifold, we have*

$$W_q^{\text{LF}} = \tilde{W}_{\check{q}(q)}^{\text{HV}}$$

where $\tilde{W}^{\text{HV}}$ is the Hori-Vafa superpotential in a suitable coordinates (see Equation (3.4)).

Proof. This follows from [7, Theorem 5.3]. $\square$

Remark 6.9. *In fact the proof of [7, Theorem 5.3] shows that the statements in Corollaries 6.7 and 6.8 and that of Theorem 6.5 are all equivalent to each other.*

Finally we have

Corollary 6.10 (=Theorem 1.5). *Let X be a compact semi-Fano toric manifold. Then the isomorphism (1.3) maps each normalized Seidel element $S_l^\circ \in QH^*(X, \omega_q)$ to the generator Z_l ($l = 1, \dots, m$) of the Jacobian ring $\text{Jac}(W_q^{LF})$, where Z_l are the monomials defined by Equation (2.2).*

Proof. By Corollary 6.8, since W^{LF} and $\tilde{W}^{HV}$ are equal, $QH^*(X, \omega_q) \xrightarrow{\sim} \text{Jac}(W_q^{LF})$ is the same as $QH^*(X, \omega_q) \xrightarrow{\sim} \text{Jac}(\tilde{W}_{q(q)}^{HV})$. By Proposition 3.15, each Batyrev element B_l is mapped to $(\exp g_l)Z_l$ for $l = 1, \dots, m$. Since $B_l = (\exp g_l)S_l^\circ$ by Proposition 6.1, it follows that S_l° is mapped to Z_l for $l = 1, \dots, m$. $\square$

Remark 6.11. *On the other hand, if the isomorphism (1.3) maps S_l° to Z_l for $l = 1, \dots, m$, then the elements $\tilde{B}_l \in QH^*(X, \omega_q)$ defined by $\tilde{B}_l := (1 + \delta_l)S_l^\circ$ satisfy the conditions (i), (ii) and (iii) of [20, Theorem 1.2], which states that these conditions completely characterize the Batyrev elements. So we have $\tilde{B}_l = B_l$ in $QH^*(X, \omega_q)$. Theorem 6.5 then follows from Proposition 6.1. Hence Corollary 6.10 is in fact also equivalent to Theorem 6.5.*

We conjecture that Corollary 6.10 holds true for *any* compact toric manifold:

Conjecture 6.12. *Let X be a compact toric manifold, not necessarily semi-Fano. Then the isomorphism (1.3) maps the normalized Seidel elements $S_l^\circ \in QH^*(X, \omega_q)$ to the generators⁷ Z_l of the Jacobian ring $\text{Jac}(W_q^{LF})$, where Z_l are monomials defined by Equation (2.2).*

Example 6.13. *Consider the semi-Fano toric surface X whose moment map image is shown in Figure 3. The disc potential W_q^{LF} and generating functions δ_i of X were computed in [5]. The key result is $n_1(\beta) = 1$ if β is an admissible disc class, and $n_1(\beta) = 0$ otherwise. Admissibility is a combinatoric condition which is easy to check, and the readers are referred to [5] for the detailed definition and results.*




FIGURE 3. An example of semi-Fano toric surface, which was denoted as X_8 in Appendix A of [5].

⁷In general W_q^{LF} is a Laurent series over the Novikov ring in contrast to Definition 2.4 for semi-Fano case. Nevertheless we can still define the monomials Z_l by Equation (2.2).

The generating functions corresponding to D_1, D_2, D_3 are

$$\begin{aligned}\delta_1(q) &= q_1 + q_1q_2 + q_1q_2q_3, \\ \delta_2(q) &= q_2 + q_1q_2 + q_2q_3 + q_1q_2q_3 + q_1q_2^2q_3, \\ \delta_3(q) &= q_3 + q_2q_3 + q_1q_2q_3\end{aligned}$$

respectively, where q_i 's are the Kähler parameters of D_i 's for $i = 1, 2, 3$. Each term in the above generation functions corresponds to an admissible disc class.

On the other hand, the mirror map is given by

$$\begin{aligned}q_1 &= \check{q}_1 \exp(2g_1(\check{q}_1, \check{q}_2, \check{q}_3) - g_2(\check{q}_1, \check{q}_2, \check{q}_3)); \\ q_2 &= \check{q}_2 \exp(-g_1(\check{q}_1, \check{q}_2, \check{q}_3) + 2g_2(\check{q}_1, \check{q}_2, \check{q}_3) - g_3(\check{q}_1, \check{q}_2, \check{q}_3)); \\ q_3 &= \check{q}_3 \exp(-g_2(\check{q}_1, \check{q}_2, \check{q}_3) + 2g_3(\check{q}_1, \check{q}_2, \check{q}_3))\end{aligned}$$

where

$$\begin{aligned}g_1(\check{q}_1, \check{q}_2, \check{q}_3) &= \sum_{(a,b,c)} \check{q}_1^a \check{q}_2^b \check{q}_3^c \frac{(-1)^{2a-b}(2a-b-1)!}{a!c!(a-2b+c)!(b-2c)!}; \\ g_2(\check{q}_1, \check{q}_2, \check{q}_3) &= \sum_{(a,b,c)} \check{q}_1^a \check{q}_2^b \check{q}_3^c \frac{(-1)^{2b-a-c}(2b-a-c-1)!}{a!c!(b-2a)!(b-2c)!}; \\ g_3(\check{q}_1, \check{q}_2, \check{q}_3) &= \sum_{(a,b,c)} \check{q}_1^a \check{q}_2^b \check{q}_3^c \frac{(-1)^{2c-b}(2c-b-1)!}{a!c!(a-2b+c)!(b-2a)!}\end{aligned}$$

where the summations are over all $(a, b, c) \in \mathbb{Z}^3$ such that the term before each factorial sign is non-negative. By Theorem 6.5, we have $1 + \delta_i(q(\check{q})) = \exp g_i(\check{q})$ for $i = 1, 2, 3$. This produces non-trivial identities between hypergeometric series, and hence a closed formula for the inverse mirror map $\check{q}(q)$:

$$\begin{aligned}\check{q}_1 &= q_1 \cdot \frac{1 + q_2 + q_1q_2 + q_2q_3 + q_1q_2q_3 + q_1q_2^2q_3}{(1 + q_1 + q_1q_2 + q_1q_2q_3)^2}; \\ \check{q}_2 &= q_2 \cdot \frac{(1 + q_1 + q_1q_2 + q_1q_2q_3)(1 + q_3 + q_2q_3 + q_1q_2q_3)}{(1 + q_2 + q_1q_2 + q_2q_3 + q_1q_2q_3 + q_1q_2^2q_3)^2}; \\ \check{q}_3 &= q_3 \cdot \frac{1 + q_2 + q_1q_2 + q_2q_3 + q_1q_2q_3 + q_1q_2^2q_3}{(1 + q_3 + q_2q_3 + q_1q_2q_3)^2}.\end{aligned}$$

REFERENCES

1. D. Auroux, *Mirror symmetry and T-duality in the complement of an anticanonical divisor*, J. Gökova Geom. Topol. GGT **1** (2007), 51–91. MR 2386535 (2009f:53141)
2. ———, *Special Lagrangian fibrations, wall-crossing, and mirror symmetry*, Surveys in differential geometry. Vol. XIII. Geometry, analysis, and algebraic geometry: forty years of the Journal of Differential Geometry, Surv. Differ. Geom., vol. 13, Int. Press, Somerville, MA, 2009, pp. 1–47. MR 2537081 (2010j:53181)
3. V. Batyrev, *Quantum cohomology rings of toric manifolds*, Astérisque (1993), no. 218, 9–34, Journées de Géométrie Algébrique d'Orsay (Orsay, 1992). MR 1265307 (95b:32034)

4. K. Chan, *A formula equating open and closed Gromov-Witten invariants and its applications to mirror symmetry*, Pacific J. Math. **254** (2011), no. 2, 275–293. MR 2900016
5. K. Chan and S.-C. Lau, *Open Gromov-Witten invariants and superpotentials for semi-Fano toric surfaces*, preprint, arXiv:1010.5287.
6. K. Chan, S.-C. Lau, and N.C. Leung, *SYZ mirror symmetry for toric Calabi-Yau manifolds*, J. Differential Geom. **90** (2012), no. 2, 177–250. MR 2899874
7. K. Chan, S.-C. Lau, N.C. Leung, and H.-H. Tseng, *Open Gromov-Witten invariants and mirror maps for semi-Fano toric manifolds*, preprint, arXiv:1112.0388.
8. K. Chan, S.-C. Lau, and H.-H. Tseng, *Enumerative meaning of mirror maps for toric Calabi-Yau manifolds*, preprint, arXiv:1110.4439.
9. C.-H. Cho, *Products of Floer cohomology of torus fibers in toric Fano manifolds*, Comm. Math. Phys. **260** (2005), no. 3, 613–640. MR 2183959 (2006h:53094)
10. C.-H. Cho and Y.-G. Oh, *Floer cohomology and disc instantons of Lagrangian torus fibers in Fano toric manifolds*, Asian J. Math. **10** (2006), no. 4, 773–814. MR 2282365 (2007k:53150)
11. K. Fukaya, Y.-G. Oh, H. Ohta, and K. Ono, *Lagrangian Floer theory and mirror symmetry on compact toric manifolds*, preprint, arXiv:1009.1648.
12. ———, *Lagrangian intersection Floer theory: anomaly and obstruction. Part I*, AMS/IP Studies in Advanced Mathematics, vol. 46, American Mathematical Society, Providence, RI, 2009. MR 2553465 (2011c:53217)
13. ———, *Lagrangian intersection Floer theory: anomaly and obstruction. Part II*, AMS/IP Studies in Advanced Mathematics, vol. 46, American Mathematical Society, Providence, RI, 2009. MR 2548482 (2011c:53218)
14. ———, *Lagrangian Floer theory on compact toric manifolds. I*, Duke Math. J. **151** (2010), no. 1, 23–174. MR 2573826 (2011d:53220)
15. ———, *Lagrangian Floer theory on compact toric manifolds II: bulk deformations*, Selecta Math. (N.S.) **17** (2011), no. 3, 609–711. MR 2827178
16. ———, *Toric degeneration and non-displaceable Lagrangian tori in $S^2 \times S^2$* , Int. Math. Res. Not. IMRN **2012** (2012), no. 13, 2942–2993, arXiv:1002.1660.
17. W. Fulton, *Introduction to toric varieties*, Annals of Mathematics Studies, vol. 131, Princeton University Press, Princeton, NJ, 1993, The William H. Roever Lectures in Geometry. MR 1234037 (94g:14028)
18. A. Givental, *A mirror theorem for toric complete intersections*, Topological field theory, primitive forms and related topics (Kyoto, 1996), Progr. Math., vol. 160, Birkhäuser Boston, Boston, MA, 1998, pp. 141–175. MR 1653024 (2000a:14063)
19. E. González and H. Iritani, in preparation.
20. ———, *Seidel elements and mirror transformations*, Selecta Math. (N.S.) **18** (2012), no. 3, 557–590, arXiv:1103.4171.
21. K. Hori and C. Vafa, *Mirror symmetry*, preprint, arXiv:hep-th/0002222.
22. H. Iritani, *An integral structure in quantum cohomology and mirror symmetry for toric orbifolds*, Adv. Math. **222** (2009), no. 3, 1016–1079. MR 2553377 (2010j:53182)
23. S.-C. Lau, N.C. Leung, and B. Wu, *A relation for Gromov-Witten invariants of local Calabi-Yau threefolds*, Math. Res. Lett. **18** (2011), no. 5, 943–956. MR 2875867
24. ———, *Mirror maps equal SYZ maps for toric Calabi-Yau surfaces*, Bull. London Math. Soc. **44** (2012), 255–270.
25. W. Lerche and P. Mayr, *On $\mathcal{N} = 1$ mirror symmetry for open Type II strings*, preprint, arXiv:hep-th/0111113.
26. W. Lerche, P. Mayr, and N. Warner, *$\mathcal{N} = 1$ special geometry, mixed Hodge variations and toric geometry*, preprint, arXiv:hep-th/0208039.
27. B. Lian, K. Liu, and S.-T. Yau, *Mirror principle. III*, Asian J. Math. **3** (1999), no. 4, 771–800. MR 1797578 (2002g:14080)
28. P. Mayr, *Summing up open string instantons and $\mathcal{N} = 1$ string amplitudes*, preprint, arXiv:hep-th/0203237.
29. ———, *$\mathcal{N} = 1$ mirror symmetry and open/closed string duality*, Adv. Theor. Math. Phys. **5** (2001), no. 2, 213–242. MR 1909365 (2003e:32028)

- 30. D. McDuff, *Quantum homology of fibrations over S^2* , Internat. J. Math. **11** (2000), no. 5, 665–721. MR 1780735 (2001i:53157)
- 31. D. McDuff and S. Tolman, *Topological properties of Hamiltonian circle actions*, IMRP Int. Math. Res. Pap. (2006), 72826, 1–77. MR 2210662 (2007e:53115)
- 32. P. Seidel, π_1 of symplectic automorphism groups and invertibles in quantum homology rings, Geom. Funct. Anal. **7** (1997), no. 6, 1046–1095. MR 1487754 (99b:57068)

DEPARTMENT OF MATHEMATICS, THE CHINESE UNIVERSITY OF HONG KONG, SHATIN, HONG KONG

E-mail address: kwchan@math.cuhk.edu.hk

DEPARTMENT OF MATHEMATICS, HARVARD UNIVERSITY, ONE OXFORD STREET, CAMBRIDGE, MA 02138, USA

E-mail address: s.lau@math.harvard.edu

THE INSTITUTE OF MATHEMATICAL SCIENCES AND DEPARTMENT OF MATHEMATICS, THE CHINESE UNIVERSITY OF HONG KONG, SHATIN, HONG KONG

E-mail address: leung@math.cuhk.edu.hk

DEPARTMENT OF MATHEMATICS, OHIO STATE UNIVERSITY, 100 MATH TOWER, 231 WEST 18TH AVE., COLUMBUS, OH 43210, USA

E-mail address: hhtseng@math.ohio-state.edu